



**CALIFORNIA**  
HEALTH BENEFITS REVIEW PROGRAM

---

Analysis of Senate Bill 576  
Health Care Coverage:  
Tobacco Cessation Services

---

A Report to the 2005-2006 California Legislature  
April 16, 2005  
*Revised May 20, 2005*  
*Revised August 22, 2005*

CHBRP 05-07



Established in 2002 to implement the provisions of Assembly Bill 1996 (*California Health and Safety Code*, Section 127660, et seq.), the California Health Benefits Review Program (CHBRP) responds to requests from the State Legislature to provide independent analysis of the medical, financial, and public health impacts of proposed health insurance benefit mandates. The statute defines a health insurance benefit mandate as a requirement that a health insurer and/or managed care health plan (1) permit covered individuals to receive health care treatment or services from a particular type of health care provider; (2) offer or provide coverage for the screening, diagnosis, or treatment of a particular disease or condition; or (3) offer or provide coverage of a particular type of health care treatment or service, or of medical equipment, medical supplies, or drugs used in connection with a health care treatment or service.

A small analytic staff in the University of California's Office of the President supports a task force of faculty from several campuses of the University of California, as well as Loma Linda University, the University of Southern California, and Stanford University, to complete each analysis within a 60-day period, usually before the Legislature begins formal consideration of a mandate bill. A certified, independent actuary helps estimate the financial impacts, and a strict conflict-of-interest policy ensures that the analyses are undertaken without financial or other interests that could bias the results. A National Advisory Council, made up of experts from outside the state of California and designed to provide balanced representation among groups with an interest in health insurance benefit mandates, reviews draft studies to ensure their quality before they are transmitted to the Legislature. Each report summarizes sound scientific evidence relevant to the proposed mandate but does not make recommendations, deferring policy decision making to the Legislature. The State funds this work through a small annual assessment of health plans and insurers in California. All CHBRP reports and information about current requests from the California Legislature are available at CHBRP's Web site, [www.chbrp.org](http://www.chbrp.org).

**A Report to the 2005-2006 California State Legislature**

**Analysis of Senate Bill 576  
Health Care Coverage: Tobacco Cessation Services**

**April 16, 2005**  
*Revised May 20, 2005*  
*Revised August 22, 2005*

**California Health Benefits Review Program  
1111 Franklin Street, 11<sup>th</sup> Floor  
Oakland, CA 94607  
Tel: 510-287-3876  
Fax: 510-987-9715  
[www.chbrp.org](http://www.chbrp.org)**

Additional free copies of this and other CHBRP bill analyses and publications may be obtained by visiting the CHBRP Web site at [www.chbrp.org](http://www.chbrp.org).

**Suggested Citation:**

California Health Benefits Review Program (CHBRP). (2005). *Analysis of Senate Bill 576 Health Care Coverage: Tobacco Cessation Services*. Report to Calif. State Legislature. Oakland, CA: CHBRP. 05-07.



## PREFACE

This report provides an analysis of the medical, financial, and public health impacts of SB 576, a bill to mandate coverage of tobacco cessation services, as introduced on February 18, 2005 and subsequently amended and transmitted to the California Health Benefits Review Program (CHBRP) in draft form on July 21, 2005. In response to this transmitted request from the California Senate Banking, Finance and Insurance Committee, CHBRP undertook this analysis pursuant to the provisions of Assembly Bill 1996 (2002) as chaptered in Section 127600, et seq. of the California Health and Safety Code. This analysis is similar to the Reports to the California State Legislature submitted on April 16, and May 20, 2005. The April 16, 2005 CHBRP analysis assessed the impacts of SB 576 as introduced on February 18, 2005. The May 20, 2005 CHBRP analysis assessed the impacts of the version of SB 576 as amended on May 3, 2005 and revised the cost estimates for Medi-Cal to appropriately reflect that Medi-Cal currently covers tobacco cessation treatment services. This analysis incorporates the July 21, 2005 draft amendments which would allow copayments for over-the-counter drugs for the treatment of tobacco cessation. It also revises the cost and public health estimates to correctly reflect the expected reduction of heart attacks as a result of the mandate.

Wade Aubry, MD, Patricia Franks, BA, Noelle Lee, BA, Harold S. Luft, PhD, and Edward Yelin, PhD, all of the University of California, San Francisco, prepared the medical effectiveness analysis. Min-Lin Fang, MLIS, of UCSF conducted the literature search. Sharon M. Hall, PhD\*, University of California, San Francisco, provided technical assistance with the literature review and expertise for the medical effectiveness analysis. Yali Bair, PhD, Richard Kravitz, MD, Janet Keyzer, RN-C, MPA, and Christina Kuenneth, MPH, all of the University of California, Davis prepared the public health impact analysis. Gerald Kominski, PhD, and Nadereh Pourat, PhD, both of the University of California, Los Angeles, prepared the analysis of the cost impact. Robert Cosway, FSA, MAAA, and Chris Girod, FSA, MAAA, both of Milliman, provided actuarial analysis. Susan Philip, MPP, and Robert O'Reilly, BS, of CHBRP staff prepared the background section and contributed to integrating the individual sections into a single report. Other contributors include Sachin Kumar, BA, Cynthia Robinson, MPP, both of CHBRP staff, and Cherie Wilkerson, who provided editing services. In addition, a subcommittee of CHBRP's National Advisory Council (see final pages of this report) reviewed the analysis for its accuracy, completeness, clarity, and responsiveness to the Legislature's request.

Jay Ripps, FSA, MAAA of Milliman recused himself from contributing to this and all other CHBRP analyses beginning March 1, 2005. His recusal is valid through his duration as acting chief actuary at Blue Shield of California.

CHBRP gratefully acknowledges all of these contributions but assumes full responsibility for all of the report and its contents. Please direct any questions concerning this report to CHBRP:

**California Health Benefits Review Program**  
**1111 Franklin Street, 11<sup>th</sup> Floor**  
**Oakland, CA 94607**  
**Tel: 510-287-3876**  
**Fax: 510-987-9715**  
[www.chbrp.org](http://www.chbrp.org)

All CHBRP bill analyses and other publications are available on CHBRP's Web site, [www.chbrp.org](http://www.chbrp.org).  
Michael E. Gluck, PhD, Director

*\*Dr. Hall is a current member of the Society for Research on Nicotine & Tobacco (SRNT). Among other things, SRNT supports assurance that tobacco cessation treatments have been integrated into regular health care delivery systems for all patients; Dr. Hall has a current research grant funded by Pfizer on a study for a smoking cessation aid, although her role is limited to advice about treatment and data collection; Dr. Hall did not declare any proprietary interest in Pfizer.*



## TABLE OF CONTENTS

EXECUTIVE SUMMARY .....	5
INTRODUCTION .....	11
I. MEDICAL EFFECTIVENESS .....	12
II. UTILIZATION, COST, AND COVERAGE IMPACTS.....	18
Present Baseline Cost and Coverage.....	18
Impacts of Mandated Coverage.....	21
III. PUBLIC HEALTH IMPACTS .....	23
APPENDICES .....	33
REFERENCES .....	57



## EXECUTIVE SUMMARY

### California Health Benefits Review Program Analysis of Senate Bill 576

This report provides an analysis of the medical, financial, and public health impacts of SB 576, a bill to mandate coverage of tobacco cessation services, as introduced on February 18, 2005 and subsequently amended and transmitted to the California Health Benefits Review Program (CHBRP) in draft form on July 21, 2005. In response to this transmitted request from the California Senate Banking, Finance and Insurance Committee, CHBRP undertook this analysis pursuant to the provisions of Assembly Bill 1996 (2002) as chaptered in Section 127600, et seq. of the California Health and Safety Code. This analysis is similar to the Reports to the California State Legislature submitted on April 16, and May 20, 2005. The April 16, 2005 CHBRP analysis assessed the impacts of SB 576 as introduced on February 18, 2005. The May 20, 2005 CHBRP analysis assessed the impacts of the version of SB 576 as amended on May 3, 2005 and revised the cost estimates for Medi-Cal to appropriately reflect that Medi-Cal currently covers tobacco cessation treatment services. This analysis incorporates the July 21, 2005 draft amendments which would allow copayments for over-the-counter drugs for the treatment of tobacco cessation. It also revises the cost and public health estimates to correctly reflect the expected reduction of heart attacks as a result of the mandate.

Senate Bill (SB) 576 would require health care service plans and insurers that provide outpatient prescription drug benefits coverage to cover the following tobacco cessation services:

- Personal counseling, which may include telephone or individual counseling.
- Prescription and over-the-counter tobacco cessation medications approved by the Food and Drug Administration.
- Benefits for tobacco cessation must comply with the Public Health Service–sponsored 2000 clinical practice guideline, “A Clinical Practice Guideline for Treating Tobacco Use and Dependence” (PHS, 2000) or its successors.

Health plans and insurers would be permitted to limit the courses of treatment or how many “quit” attempts the member makes, to two per every twelve-month period.

SB 576 would allow for copayments for over-the-counter (OTC) and prescription drugs but they are not to exceed \$15 copayment.<sup>1</sup>

SB 576 would apply to health care services plans licensed by Knox-Keene<sup>2</sup> and to health insurance policies regulated under the California Insurance code.

#### I. Medical Effectiveness

##### *Outcomes*

- Current evidence-based systematic reviews and meta-analyses all conclude that counseling interventions, brief advice from physicians and clinical staff, and Food and Drug

---

<sup>1</sup> SB 576 would add §1367.27 of the California Health and Safety Code and §10123.175 of the California Insurance Code. SB 576 excludes specialized health care service plans, such as plans that provide a specific type of service such as vision or dental.

<sup>2</sup> Health maintenance organizations in California are licensed under the Knox-Keene Health Care Services Plan Act, which is part of the California Health and Safety Code.



Administration (FDA)-approved pharmacotherapy are effective treatments for tobacco cessation, as measured by abstinence.

- The 2000 U.S. Public Health Service Clinical Guideline (PHS, 2000) determined that there is a strong dose-response relation between the session length of person-to-person contact and successful treatment outcomes. Evidence indicates that a minimum of four counseling sessions of at least 30 minutes each, including proactive telephone counseling and individual counseling, is effective.
- Brief advice from a physician and clinical staff is also an effective intervention for tobacco cessation. The evidence indicates that physician advice to quit tobacco use increases abstinence rates.
- Prescription and over-the-counter nicotine replacement medications increase abstinence rates, as does bupropion, a non-nicotine prescription drug. Medications can double the likelihood that a patient will cease tobacco use.
- Combinations of interventions have incremental increases in effectiveness.
- Full coverage of tobacco cessation benefits significantly increased quit rates in one trial. In a randomized controlled trial, compliance with the U.S. Public Health Service Clinical Guideline (PHS, 2000) resulted in higher abstinence rates compared with the control group.

*Caveat*

- The basic conclusion about the effectiveness of interventions for tobacco cessation is not likely to be altered or diminished with the publication of new studies.

## **II. Utilization, Cost, and Coverage Impacts**

- 20,368,000 individuals are currently enrolled in health plans regulated by the Knox-Keene Act or insured by policies regulated under the California Insurance Code. Currently, 95% of this population have coverage for prescription drugs and would be affected by SB 576—this includes 12,924,000 adults ages 18 and older.
- 7,052,000 insured adults (51.6%) currently have coverage for tobacco cessation benefits, though those covered benefits generally do *not* include coverage for over-the-counter (OTC) medications except for those covered by MediCal managed care organizations. Coverage tends to include physician visits, counseling, and prescription medication.
- Total health expenditures are projected to increase by \$77,098,000 (0.128%). This overall increase includes a net savings of \$11,545,000 that is related to the reduced out-of-pocket expenditures for non-covered services and a savings of \$3,418,000 that is related to lower health care use among additional smokers who would quit as a result of access to tobacco



cessation services under the mandate<sup>3</sup>. These savings represent the short-term (i.e., one-year) savings resulting from reduced use of health care services among those who quit smoking. The potential long-term savings of quitting are likely to be substantial due to reductions in the rate of smoking-related illnesses. However, calculations of such savings are considered beyond the scope of this analysis.

- The per member per month (PMPM) increases in health expenditures and premiums vary by covered population, due to variations in the current coverage levels of smoking cessation benefits and other factors. The expected increase in PMPM expenditures for large group health maintenance organizations (HMO) members is 0.120% or \$ 0.31. Values for other populations are summarized in Table 10.

---

<sup>3</sup> The cost savings calculation incorporates savings associated with reduced hospitalization for acute myocardial infarction (AMI) and savings associated with fewer low-birth weight deliveries. This analysis revises the cost savings to reflect the correct reduction in expected AMIs.



**Table 1. Summary of Coverage, Utilization, and Cost Effects of SB 576**

<b>Total Insured Population = 20,368,000</b>	<b>Before Mandate</b>	<b>After Mandate</b>	<b>Increase/Decrease</b>	<b>% Change After Mandate</b>
<b><u>Coverage</u></b>				
Percentage of insured individuals with coverage for mandated benefit	55.1%*	95.0%	39.9%	72.4%
Number of insured individuals in California with coverage for the benefit	11,217,000*	19,341,000	8,124,000	72.4%
Percentage of insured individuals 18 years and older with coverage for mandated benefit	51.6%*	94.6%	43.0%	83.3%
Number of insured individuals 18 years and older in California with coverage for the benefit	7,052,000*	12,924,000	5,872,000	83.3%
<b><u>Total Unit Cost</u></b>				
Counseling	\$187.50	\$300.00	\$112.50	60.0%
Prescription Drugs	\$300.00	\$300.00	—	0%
OTC Drugs	\$285.00	\$285.00	—	0%
Average cost of cessation treatment for those with a covered benefit	\$487.50	\$614.55	\$127.05	26.1%
Average cost of cessation treatment for those without a covered benefit	\$285.00	N/A		
Average savings in health expenditures during first year for smokers who quit	\$192.14	\$192.14	—	0%
<b><u>Utilization</u></b>				
Percentage members 18 years and older who smoke with covered benefit and who try tobacco cessation	10.0%	10.2%	0.2% point	2.0%
Percentage members 18 years and older who smoke without covered benefit and who try tobacco cessation	4%	N/A	N/A	N/A



**Table 1. Summary of Coverage, Utilization, and Cost Effects of SB 576 (continued)**

<b>Total Insured Population = 20,368,000</b>	<b>Before Mandate</b>	<b>After Mandate</b>	<b>Increase/Decrease</b>	<b>% Change After Mandate</b>
<b>Utilization</b>				
1-Year success rate for members 18 years and older who smoke with covered benefit and who try tobacco cessation	20%	20%	0%	0%
1-Year success rate for members 18 years and older who smoke without covered benefit and who try tobacco cessation	8%	8%	0.0%	0.0%
<b>Annual Expenditures</b>				
Premium expenditures by private employers for group insurance	35,360,055,000	35,412,130,000	52,075,000	0.147%
Premium expenditures by individuals with group insurance, CalPERS, or Healthy Families	10,261,105,000	10,276,255,000	15,150,000	0.148%
Premium expenditures for individually purchased insurance	3,818,726,000	3,831,714,000	12,988,000	0.340%
CalPERS employer expenditures	2,212,881,000	2,214,219,000	1,338,000	0.060%
Medi-Cal state expenditures	3,939,663,000	3,939,663,000	0	0.000%
Healthy Families state expenditures	347,858,000	347,928,000	70,000	0.020%
Member copayments	4,074,893,000	4,081,915,000	7,022,000	0.172%
Expenditures for non-covered services	11,545,000	-	11,545,000	-100.000%
<b>Total annual expenditures</b>	<b>60,026,726,000</b>	<b>60,103,824,000</b>	<b>77,098,000</b>	<b>0.128%</b>

Source: California Health Benefits Review Program, 2005.

Notes: \* These members have coverage for tobacco cessation benefits, though those covered benefits generally do not include coverage for over-the-counter (OTC) medications except for those covered by MediCal managed care organizations.

The population includes individuals and dependents in California who have private insurance (group and individual) or are enrolled in public plans subject to the Health and Safety Code, including CalPERS, Medi-Cal, or Healthy Families.

All population figures include enrollees aged 0-64 years, except the Medi-Cal population, which includes dually eligible Medicare/Medi-Cal recipients of all ages. Employees and their dependents who receive their coverage from self-insured firms are excluded because these plans are not subject to mandates.

Key: CalPERS = California Public Employees' Retirement System.



### III. Public Health Impacts

- Tobacco-related disease imposes a significant cost impact on individuals, employers, and society in terms of illness, medical costs, and loss of life and productivity.
- Tobacco cessation services have been proven effective in reducing the short-term disease and cost impacts of low birth weight deliveries and acute myocardial infarction within the first year after cessation.
- Tobacco cessation services have the potential to result in long-term public health benefits and cost savings by reducing exposure to the multiple mechanisms by which smoking causes disease, disability, and death.
- During the first year after implementation, this mandate is estimated to reduce, statewide, low birth weight deliveries by approximately 58 cases and would reduce acute myocardial infarction by 12 cases.<sup>4</sup>

---

<sup>4</sup> This analysis revises the public health impact estimate to reflect the correct reduction in expected AMIs by basing the estimate on the corrected incidence rate of AMI which is 0.219 (as opposed to 2.19 as stated in prior CHBRP analyses of SB 576).



## INTRODUCTION

Tobacco use is the leading cause of preventable death in the United States and is associated with a variety of diseases. The 2004 U.S. Surgeon General’s Report concluded that tobacco cessation has immediate and long-term benefits by reducing risks for diseases caused by smoking and improving health in general (DHHS, 2004).

Senate Bill (SB) 576 would require health care service plans and insurers that provide outpatient prescription drug benefits coverage to cover the following tobacco cessation services:

- Personal counseling, which may include telephone or individual counseling.
- Prescription and over-the-counter tobacco cessation medications approved by the Food and Drug Administration.

Benefits for tobacco cessation must comply with the Public Health Service–sponsored 2000 clinical practice guideline, “A Clinical Practice Guideline for Treating Tobacco Use and Dependence” (PHS, 2000) or its successors.

Health plans and insurers would be permitted to limit the courses of treatment or how many “quit” attempts the member makes, to two per every twelve-month period.

SB 576 would allow for copayments for over-the-counter (OTC) or prescription drugs but they are not to exceed \$15 copayment. The bill is silent on copayments for all other services. Therefore, this analysis assumes that the usual copayment levels for counseling and physician office visits would continue to apply.<sup>5</sup>

Enrollees and their providers may decide at the beginning of a tobacco cessation course what services and products they prefer in order to quit. Coverage for tobacco cessation services, including OTC drugs, would only “be available upon the order of an authorized provider.”

SB 576 would apply to health care services plans licensed by Knox-Keene<sup>6</sup> and to health insurance policies regulated under the California Insurance code.

Currently no other state requires insurers to cover tobacco cessation programs as a health benefit. Many legislatures have considered such legislation, including New York, Maryland, Wisconsin, and Oklahoma.

Over the past several years, California has taken measures to decrease the number of smokers and prevent an increase in the number of new smokers. For example, Californians have had increased access to tobacco cessation education and services through the California Department of Health Services Tobacco Control Section. The California Tobacco Tax and Health Promotion Act of 1988 (Proposition 99) increased the state surtax on cigarettes and other tobacco-related products. Revenues from the “tobacco tax” were appropriated for tobacco-related research, tobacco cessation efforts, and health education and

---

<sup>5</sup> SB 576 would add §1367.27 of the California Health and Safety Code and §10123.175 of the California Insurance Code. SB 576 excludes specialized health care service plans, such as plans that provide a specific type of service such as vision or dental.

<sup>6</sup> Health maintenance organizations in California are licensed under the Knox-Keene Health Care Services Plan Act, which is part of the California Health and Safety Code.



health care for medically-indigent families. In 1995, California enacted a smoke-free workplace law in an effort to reduce the public health burden of second-hand smoke inhalation.

Tobacco settlement monies provided California with approximately \$1 billion a year. However, beginning with the 2002-2003 budget, the state began to divert its share of tobacco settlement fund revenues (TFR) from health programs to debt repayment (Legislative Analyst's Office, 2002). Since 2003, the state has continued to divert all the TFR toward debt repayment.<sup>7</sup>

## **I. MEDICAL EFFECTIVENESS**

This medical effectiveness analysis reviews the literature on the effectiveness of counseling, brief cessation interventions, and medications approved by the Federal Drug Administration (FDA) for the use of tobacco cessation treatment. Counseling interventions include counseling in person or via telephone and in individual or group therapy. Counseling is often provided by physicians, nurses, peer counselors, social workers, psychologists, or psychiatrists. Pharmacotherapy includes nicotine replacement therapy (NRT), which can be administered by chewing gum, nasal spray, transdermal patch, inhaler, or lozenge. Bupropion SR, which does not contain nicotine, has been FDA-approved as a tobacco cessation aid for people who smoke 10 or more cigarettes daily and are at least 18 years old. SB 576 does not specify if interventions are directed at adolescents or adults; thus, the medical effectiveness review does not exclude any population.

Extensive literature has been published on the effectiveness of various tobacco cessation interventions. Most studies reviewed employed abstinence as the primary outcome measure in evaluating the effectiveness of various interventions. Although continuous abstinence is desirable, varying definitions of relapse create difficulty in evaluating prolonged-abstinence rates in patients. However, because most relapses occur within the first three months after tobacco cessation, many systematic reviews of the literature included only those studies with follow-up of at least five months (PHS, 2000). Thus, this medical effectiveness analysis considers interventions that result in tobacco abstinence for at least five months to be effective.

The results of the review of the scientific literature relevant to SB 576 are divided into three major categories of interventions: (1) counseling, (2) brief advice given by physicians and clinical staff, and (3) pharmacotherapies. Although other interventions for tobacco cessation may exist and are effective, this review only discusses interventions proposed in the mandate.

The scope of the literature search included the following outcomes:

- Tobacco cessation (quit smoking for less than six months, quit smoking for six months to one year, quit smoking for more than one year, quit smoking for five years, quit smoking for ten years)
- Recidivism rate (smoking not resumed for the above time intervals)
- Improved pulmonary function tests (forced expiratory volume at 1 second [FEV1]/forced vital capacity [FVC])

---

<sup>7</sup> Personal communication with member of the State of California Tobacco Education and Research Oversight Committee, March 28, 2005



However, most reviews and meta-analyses considered abstinence from tobacco use for at least five months the primary outcome measure. The literature search was conducted through PubMed and the Cochrane Library for relevant research published between 2000 and 2005 and was limited to English abstracts. Internet searches were conducted and focused on several Websites, including National Commission on Quality Assurance (Health Plan Employer Data and Information Set measures), Centers for Disease Control and Prevention, U.S. Preventive Services Task Force (USPSTF), American Cancer Society, American Cancer Institute, American Lung Association, and the Task Force on Community Preventive Services. A description of methods used to conduct the medical effectiveness review, and the process used to “grade” the evidence of effectiveness can be found in Appendix A: Literature Review Methods. Summary tables with detailed findings from the literature can be found in Appendix B: Summary of Findings on Medical Effectiveness of Tobacco Cessation Interventions.

### **Evidence-Based Reviews and Meta-analyses on Tobacco Cessation Interventions**

Much literature has been published on the effectiveness of various tobacco cessation interventions. Thousands of clinical trials have been published; consequently, this analysis does not review the results of individual trials, but rather reviews the recommendations and conclusions of evidence-based reviews and meta-analyses. Meta-analyses published in the Cochrane Library undergo rigorous review, and many meta-analyses were updated in 2004. (Results of these reviews are listed in Appendix B, Table B-1). In addition, the U.S. Public Health Service conducted a thorough review of articles published on the effectiveness of tobacco cessation interventions, despite the variety of interventions and the number of variables (i.e., length of counseling sessions, individual versus group therapy, etc).

In 2000, the U.S. Public Health Service released *Treating Tobacco Use and Dependence: Clinical Practice Guideline* based on the results of a systematic review and meta-analysis of the existing scientific literature (PHS, 2000). The 2000 guideline updates a previous guideline, *Smoking Cessation, Clinical Practice Guideline No. 18*, published in 1996 by the Agency for Health Care Policy and Research (now the Agency for Healthcare Research and Quality [AHRQ]), U.S. Department of Health and Human Services. The new guideline, which reviews evidence published through January 1, 1999, was written in response to new, effective clinical treatments for tobacco dependence that had been identified since the 1996 guideline was published. The following seven Federal government and nonprofit organizations sponsored the updated guideline: AHRQ, the Centers for Disease Control and Prevention (CDC), National Cancer Institute (NCI), National Heart, Lung, and Blood Institute (NHLBI), National Institute on Drug Abuse (NIDA), Robert Wood Johnson Foundation (RWJF), and University of Wisconsin Medical School’s Center for Tobacco Research and Intervention (CTRI).

The key conclusions of the 2000 U.S. Public Health Service Clinical Practice Guideline (PHS, 2000), based on scientific evidence, are as follows:

1. Tobacco dependence is a chronic condition that often requires repeated intervention. However, effective treatments exist that can produce long-term or even permanent abstinence.
2. Effective tobacco dependence treatments are available.
3. Brief tobacco dependence treatment is effective.



4. There is a strong dose-response relation between the intensity of tobacco dependence counseling and its effectiveness. Treatments involving person-to-person contact (via individual, group, or proactive telephone counseling) are consistently effective, and their effectiveness increases with treatment intensity (e.g., minutes of contact).
5. Three types of counseling and behavioral therapies were found to be especially effective:
  - Provision of practical counseling (problem solving/skills training);
  - Provision of social support as part of treatment (intra-treatment social support); and
  - Help in securing social support outside of treatment (extra-treatment social support).
6. Numerous effective pharmacotherapies for tobacco cessation now exist.
  - Five first-line pharmacotherapies were identified that reliably increase long-term tobacco abstinence rates:
    - Bupropion SR
    - Nicotine gum
    - Nicotine inhaler
    - Nicotine nasal spray
    - Nicotine patch
  - Two second-line pharmacotherapies were identified as efficacious and may be considered by clinicians if first-line pharmacotherapies are not effective:
    - Clonidine
    - Nortriptyline
  - OTC nicotine patches are effective relative to placebo, and their use should be encouraged.
7. Tobacco dependence treatments are both clinically effective and cost effective relative to other medical and disease prevention interventions.

Compared with the 1996 Clinical Practice Guideline, the 2000 U.S. Public Health Service Clinical Practice Guideline (PHS, 2000) concluded that there was stronger evidence of the association between counseling intensity and successful treatment outcomes. The updated Clinical Practice Guideline disclosed evidence for additional efficacious counseling strategies and for more efficacious pharmacologic treatment strategies. The Clinical Practice Guideline indicated strong evidence that tobacco cessation benefits that cover at least four counseling sessions of at least 30 minutes each, including proactive telephone counseling and individual counseling, are effective. The report notes that person-to-person treatment delivered for four or more sessions appears especially effective in increasing abstinence rates. There is a clear trend for abstinence rates to increase with total contact time, with 31–90 minutes producing abstinence rates that were significantly higher than were rates produced by 1–3 minutes of total contact time. The estimated odds ratio was highest for total contact time ranging from 91–300 minutes. Although the U.S. Public Health Service–sponsored Clinical Practice Guideline concluded that various interventions were effective in tobacco cessation, results of the meta-analysis conducted for the guideline indicate that some interventions may be more effective than others.



**Table 2. Efficacy of and Estimated Abstinence Rates for Various Types of Formats (n = 58 Studies)**

<b>Format</b>	<b>Number of Arms<sup>1</sup></b>	<b>Estimated Odds Ratio</b>
No format	20	1.0
Self-help	93	1.2
Proactive telephone counseling	26	1.2
Group counseling	52	1.3
Individual counseling	67	1.7

Source: U.S. Public Health Service, 2000.

Notes: (1) An arm can be defined as any of the treatment groups in a randomized clinical trial. Most randomized trials have two "arms," but some have three or more.

The Centers for Medicare and Medicaid Services (CMS) published *Decision Memo for Smoking and Tobacco Use Cessation Counseling* in March 2005 (CMS, 2005) and considered the recommendations of the 2000 guideline (PHS, 2000). CMS determined that "...the evidence is adequate to conclude that smoking and tobacco use cessation counseling, based on the current U.S. Public Health Service (PHS) Guideline, is reasonable and necessary for a patient with a disease or adverse health effect" that has been linked to tobacco use. CMS also had commissioned the RAND Corporation to conduct an evidence-based systematic review of which interventions are most effective for encouraging older smokers to quit (DHHS, 2000).

Major conclusions from the RAND report include:

1. Individual, telephone, and group counseling are all effective, with individual counseling being possibly the most effective.
2. There is consistent evidence from multiple analyses that greater intensity of counseling yields higher tobacco cessation rates.
3. NRT, clonidine, and bupropion are all effective as pharmacotherapy for tobacco cessation, although clonidine is not approved by the FDA for this use.
4. There is good evidence that both medical and non-medical providers are effective at delivering tobacco cessation services, but conflicting evidence about the relative degree of effectiveness between provider types exists.
5. Interventions with follow-up calls or visits are more effective than those without.

The authors of the RAND report conducted a meta-regression<sup>8</sup> analysis comparing an intervention with a control group to determine effectiveness by provider. The data show that many different types of providers are effective, but the trend indicates that physicians may be the most effective intervention providers.

<sup>8</sup> A meta-regression analysis aims to relate the size of effect to one or more characteristics of the studies involved.



**Table 3. Relative Provider Effectiveness**

<b>Provider</b>	<b>Adjusted Odds Ratio</b>	<b>95% Confidence Interval</b>
Physician	3.02	2.62-3.48
Psychiatrist/psychologist	2.68	1.79-4.00
Nurse	2.38	1.87-3.03
Counselor	1.87	1.35-2.61
Unknown	1.41	1.09-1.83
Other (self-help, etc.)	1.37	1.15-1.65

Source: DHHS (2000).

However, another meta-analysis examined 30 additional studies not included in the previous 1996 and 2000 U.S. Public Health Service clinical practice guidelines and found different results. Although psychologists, physicians, and nurses were effective in delivering tobacco cessation interventions, psychologists were most effective; the odds ratio for psychologists was 1.94, and the odds ratio for physicians was 1.87 (Mojica et al., 2004). Because there are more data on some clinician types than others, evaluating which clinician types are more effective is difficult. Consequently, CMS determined that all types of clinicians appear to be effective, with no one group more effective than another (CMS, 2005).

Key findings from the RAND report are that all forms of counseling have statistically significant effects on tobacco cessation and that nicotine replacement therapy (NRT) is more effective than the control group (usually placebo) in promoting tobacco cessation.

### **Effect of Insurance Coverage and Use of the U.S. Public Health Service Clinical Practice Guideline on Tobacco Cessation**

In a randomized controlled trial, Schauffler et al. (2001) assigned the control group a self-help kit (video and pamphlet) and assigned the treatment group a self-help kit and fully covered benefits for OTC NRT gum and patch and participation in a group behavioral cessation program with no patient cost sharing. The quit rates after one year of follow up were 18% in the treatment group and 13% in the control group (odds ratio = 1.6). The authors concluded that the full coverage of a tobacco-dependence treatment benefit implemented in two Independent Practice Association (IPA) model health maintenance organizations (HMOs) in California was an effective strategy for significantly increasing quit rates, quit attempts, and use of nicotine gum and patch in adult smokers.

A pilot demonstration in Washington State (Ringgen et al., 2002) implemented the federal Clinical Practice Guideline for tobacco cessation (PHS, 2000). In this observational study, union workers chose a one-call or a more intensive five-call tobacco cessation counseling plan provided by the Group Health Cooperative's Free and Clear program. Medications were limited to the nicotine patch, nicotine gum, and bupropion. The overall point prevalence quit rate for the population was 27.5% (one-call counseling, 25.5%; five-call counseling, 28.8%). The authors concluded that a combination of behavioral counseling



and medication therapy is most effective in treating tobacco dependency and that the addition of tobacco cessation programs to health benefit packages could provide valuable preventive benefits.

Katz et al. (2002) conducted a nonrandomized, controlled trial to determine the actual effectiveness of the guideline in primary care practice. The guideline-derived intervention included routine identification and brief counseling of smokers by nurses and medical assistants in addition to free NRT and telephone counseling of those smokers who were willing to make a quit attempt. The intervention was given at a family practice setting over a two-month period. After a two-month follow-up of 651 adults, significantly more intervention patients reported abstinence (21% versus 4%). At six months, the difference was not statistically significant (intervention, 21% versus control, 11%).

In another trial, Katz et al. (2004) performed a randomized, controlled trial of the effectiveness of the U.S. Public Health Service Clinical Practice Guideline (PHS, 2000) in 2,163 adult patients at eight community-based primary care clinics in southern Wisconsin. The intervention consisted of a tutorial for intake clinicians, group and individual performance feedback for intake clinicians, use of a modified vital signs stamp, an offer of free nicotine replacement therapy, and proactive telephone counseling. There was a significant difference in the percentage of patients who reported continuous abstinence at six months (intervention, 10.9% versus control, 3.8%, odds ratio = 3.4).

### **Limitations of the Analysis**

Despite the abundance of literature on various tobacco cessation interventions, there may be a publication bias. Studies with significant, favorable results may be more likely to be published than studies with neutral or insignificant results. This publication bias would affect the results of evidence-based systematic reviews and meta-analyses. In addition, although the reviews considered abstinence for five months as the primary outcome, many studies did not measure relapse rates. For example, the number of patients who remain abstinent for six months, but then resume smoking two years after the conclusion of a study is not measured.

### **Conclusions**

Evidence-based systematic reviews and meta-analyses all conclude that counseling interventions, brief advice from physicians and clinical staff, and FDA-approved pharmacotherapy are effective treatments for tobacco cessation, as measured by abstinence.

Various forms of counseling are effective. The 2000 U.S. Public Health Service Clinical Practice Guideline (PHS, 2000) determined that there is a strong dose-response relation between the session length of person-to-person contact and successful treatment outcomes. In addition, The Task Force on Community Preventive Services found strong evidence that interventions to increase cessation (a combined provider reminder *plus* provider education *with or without* patient education program; multicomponent interventions including telephone support for persons who want to stop using tobacco) are effective (CDC, 2005).

Meta-analyses have also demonstrated that brief advice from a physician and clinical staff is also an effective intervention for tobacco cessation. The 2000 U.S. Public Health Service Clinical Practice



Guideline (PHS, 2000) notes that evidence shows that physician advice to quit smoking increases abstinence rates.

Various NRTs are also effective. Prescription and OTC nicotine replacement medications increase abstinence, as does bupropion, a non-nicotine prescription drug. NRT medications include gum, dermal patches, inhalers, nasal sprays, and lozenges. For each medication, the findings of the meta-analysis conducted for the 2000 U.S. Public Health Service Clinical Practice Guideline (PHS, 2000) are similar to the effects found in more recent Cochrane reviews (see Appendix B). The 2000 Clinical Practice Guideline determined that both prescription and OTC nicotine replacement medication and bupropion are highly effective in tobacco cessation interventions. Although each intervention is effective on its own, interventions complement each other and are more effective in tobacco cessation when used together.

## **II. UTILIZATION, COST, AND COVERAGE IMPACTS**

### **Introduction**

According to CHBRP's estimates, there are 20,368,000 insured Californians currently enrolled in health plans regulated by the Knox-Keene Act or insured by policies regulated under the California Insurance Code. Currently, 95% of this population have coverage for prescription drugs and would be affected by SB 576—this includes 12,924,000 adults ages 18 and older.

According to CHBRP estimates, 9,151,000 insured Californians (44.9%) currently do not have coverage for tobacco cessation services. CHBRP's analysis assumes that insurance plans that currently cover tobacco cessation services use a combination of counseling, prescription medications, and physician contact (Javitz et al., 2004). Those with tobacco cessation are also likely to use OTC medications, such as transdermal patches and nicotine gum, to assist with their efforts (Javitz et al., 2004), although OTC medications are not currently covered by most health plans.

### **Present Baseline Cost and Coverage**

#### Current coverage of the mandated benefit (3(i))

An estimated 11,217,000 insured Californians (55.1%) currently have coverage for the mandated benefit. Smoking prevention services are particularly important in children less than 18 years of age, whereas tobacco cessation services are utilized almost exclusively by those 18 years of age and older.

CHBRP estimates that 7,052,000 insured adult Californians ages 18 and older (51.6%) currently have coverage for tobacco cessation, including:

- 1,652,000 Medi-Cal recipients in HMOs;
- 31,000 Healthy Family recipients in HMOs;
- 585,000 California Public Employees' Retirement System (CalPERS) members in HMOs;
- 528,000 persons with individually purchased coverage;
- 4,256,000 persons with employment-based coverage.



## Current utilization levels and costs of the mandated benefit (Section 3(h))

### *Current utilization*

While a majority of smokers attempt to quit in California (61.5%), only a small proportion participate in a formal cessation assistance program (see Table 5). CHBRP estimates that about 1,187,000 insured adult Californians currently smoke and have coverage for tobacco cessation services, and that on average about 10% of these current smokers annually use tobacco cessation services—specifically, participating in a formal cessation program. This assumption is based on findings from Curry et al. (1998) and discussions with an expert in the field. In addition, the utilization rate of 10% is consistent with research conducted in other states by Milliman, and an analysis by Mercer prepared for the Maryland Health Care Commission<sup>9</sup>.

### *Current average cost of tobacco cessation services*

Currently the average cost for the package of covered tobacco cessation services—including counseling and prescription drugs—is estimates to be \$487.50. This analysis assumes that advice for tobacco cessation is generally provided as part of a regular physician visit and that enrollees or members would not specifically make a doctor’s office appointment solely to discuss tobacco cessation treatment, so we estimate no additional physician costs specifically for tobacco cessation.

For insured individuals without coverage for tobacco cessation coverage, those who seek to quit use OTC medications costing on average \$285 (Levitz et al., 2004).

For the purposes of calculating cost impacts, CHBRP estimates, based on the review of the literature and discussion with an expert<sup>10</sup> in the field:

- a 20% success rate for those with covered benefits who attempt to quit.
- a 8% success rate for those without covered benefits, who attempt to quit.

### *Potential cost offsets or savings in the short-term*

Those who quit tobacco experience measurable long-term improvements in health status. Tobacco cessation also produces long-term savings in medical expenditures. Calculations of long-term savings in medical expenditures are considered beyond the scope of CHBRP analyses. However, quitting also produces short-term savings in health expenditures related to fewer inpatient stays and ambulatory care visits related to low birth-weight deliveries, and a reduction in acute myocardial infarction (AMI), or heart attacks.

**Low Birth Weight Deliveries.** Based on the assumptions described in the public health section (see page 25), CHBRP estimates the mandate could result in 58 fewer low birth weight deliveries statewide<sup>11</sup>. The average savings per avoided low birth rate delivery is estimated to be approximately \$36,000. This calculation is derived from the 1999 Lightwood study, which estimated \$21 million saved (in 1995 dollars) as a result of 1,300 fewer low birth rate deliveries. This estimated savings was then was updated

---

<sup>9</sup> On page 22 of the Maryland Health Care Commission, “Annual Mandated Health Insurance Services Evaluation” December 31, 2004 report, the study estimates 10.4% of smokers will attempt to quit using smoking cessation services (52% of smokers attempt to quit, of whom 20% would use smoking cessation services).

<sup>10</sup> Personal communication with Sharon Hall, PhD, on March 2, 2005

<sup>11</sup> This is out of the total of 32,218 insured pregnant women who were smokers prior to the mandate.



to 2005 terms at a rate of 8.4% per year.<sup>12</sup> Therefore *for each smoker who quits* as a result of the mandate, \$117 is expected to be saved for avoided low birth weight deliveries.

**Acute Myocardial Infarction.** Based on the literature described in the public health section (see page 27) CHBRP estimates the mandate could result in 12 fewer hospitalizations due to AMI<sup>13</sup>. The average savings per avoided stroke and AMI hospitalization is estimated to be approximately \$106,678. This calculation is derived from the 1997 Lightwood study (Lightwood and Glantz, 1997) which estimated an approximate \$44 million savings (1995 dollars) in one year due to reduced numbers of AMI and strokes (based on 924 fewer hospitalizations for AMI and 538 fewer hospitalizations for strokes). Using this estimate in savings, CHBRP calculated the expected total savings (for both AMI and stroke) per avoided AMI, and then was updated to 2005 terms at a rate of 8.4% per year. Therefore *for each smoker who quits* as a result of the mandate, \$75 is expected to be saved for avoided AMIs.

In total, CHBRP estimates that quitting produces an average first-year savings in health care expenditures of about \$192 per smoker who quits.

The extent to which costs resulting from lack of coverage are shifted to other payers, including both public and private entities. (Section 3(f))

CHBRP estimates no shift in costs among private or public payers as a result of current coverage. In the long-run, to the extent that smokers are more likely to require custodial nursing home services, reductions in smoking may produce reductions in nursing home expenditures under the Medi-Cal program.

Public demand for coverage (Section 3(j))

Based on the volume and extent of information received from outside parties, there is strong public interest in SB 576 (see Appendix E). Based on criteria specified under AB 1996 (2002), CHBRP is to report on the extent to which collective bargaining entities negotiate for and the extent to which self-insured plans currently have coverage for the benefits specified under the proposed mandate. Currently, the largest public self-insured plan, CalPERS preferred provider organization (PPO) plan explicitly excludes “drugs designed solely for or used to deter smoking.” Based on conversations with the largest collective bargaining agents in California, no evidence exists that unions currently include such detailed provisions during the negotiations of their health insurance policies. In order to determine whether any local unions engage in negotiations at such detail, they would need to be surveyed individually.<sup>14</sup> The Pacific Business Group on Health (PBGH) Negotiating Alliance, a large-group purchasing coalition that negotiates on behalf of large employers in California, worked with the CDC to include coverage guidelines for tobacco cessation prescription drug treatment back in 1994 and then expanded that to a comprehensive tobacco cessation approach in 2000.

---

<sup>12</sup> This trend rate was based on the average annual increase in California HMO premiums from 1995 to 2004 as measured by the Milliman Intercompany HMO Rate Survey.

<sup>13</sup> This is out of a total of 2,299,792 insured adults who were smokers prior to the mandate.

<sup>14</sup> Personal communication with SEIU and California Labor Federation on February 8, 2005



## Impacts of Mandated Coverage

### How will changes in coverage related to the mandate affect the benefit of the newly covered service and the per-unit cost? (Section 3(a))

CHBRP estimates that the unit cost of covered tobacco cessation services will increase from \$487.50 to \$614.55 (26.1%), primarily due to increased use of counseling services (see Table 1).

CHBRP estimates that the average effectiveness of tobacco cessation will increase, because those with coverage are assumed to have a 20% success rate, whereas those without coverage have only an 8% success rate. We assume this differential success rate because those without coverage would primarily rely only on OTC medications and would not have coverage for other types of interventions such as counseling.

### How will utilization change as a result of the mandate? (Section 3(b))

CHBRP estimates that the proportion of smokers currently without coverage who use tobacco cessation treatment in an attempt to quit annually will increase from 4.0% to 10.2% as a result of the mandate. Based on findings from the literature (Curry et al., 1998, Schauffler, et al., 2001, Kaper, et al, 2005), utilization is expected to increase as a result of the increased coverage for tobacco cessation treatment. In addition, among plans that already cover tobacco cessation, the proportion of smokers with coverage who attempt to quit is expected to increase as a result of the mandate, from 10.0% to 10.2%. Specifically, the mandate would limit the copayment amount for OTC products and prescription drugs to a maximum of \$15.<sup>15</sup> All other tobacco cessation services, such as physician visits, counseling sessions are not subject to copayment limits and CHBRP assumes that the usual copayment levels for these services would continue to apply.

The expected increase in utilization following the mandate is small given that the benefits and copayments requirements are similar to current smoking cessation programs. One exception is that most programs do not currently cover OTC drugs. In addition, members would be making utilization decisions based on a mutual decision between themselves and their provider about which services would be used in any given quit attempt cycle and obtain the appropriate authorizations for those services. This is expected to dampen any potential surges in utilization for any one service, since quit cycles are limited to two per year and members would be required to obtain authorization, for services, such as prescription drugs.

### To what extent does the mandate affect administrative and other expenses? (Section 3(c))

This mandate will likely increase the administrative expenses for health plans, but this increase is expected to be in proportion to the increase in health care costs. Claims administration costs may go up slightly due to an increase in claims for smoking cessation. Health plans and insurers will have to modify

---

<sup>15</sup> The draft amended version of SB 576 requested for CHBRP analysis on July 21, 2005, is silent as to the supply of drugs to which the copayment would apply. For the purposes of analysis, CHBRP assumes the copayment would apply for a one-month supply for OTC drugs and a one-month supply for prescription drugs.



some insurance contracts and member materials to reflect the new services. In addition, health plans and insurers would need to determine how to administer the tobacco cessation benefit to comply with the mandate to cover over-the-counter tobacco cessation drugs. Health plans and insurers include a component for administration and profit in their premiums. The estimated impact of this mandate on premiums includes the assumption that plans and insurers will apply their existing administration and profit loads to the marginal increase in health care costs produced by the mandate. Therefore, although there may be administrative costs associated with the mandate, administrative costs as a proportion of the premium would not change.

#### Impact of the mandate on total health care costs (Section 3(d))

SB 576 will increase total health care expenditures from \$60,026,726,000 to \$60,103,824,000 for the 20,368,000 insured individuals affected by this mandate—an increase of \$77,098,000, which equals 0.128% of total expenditures for this insured population.

#### Costs or savings for each category of insurer resulting from the benefit mandate (Section 3(e))

For each major category of payer, total annual expenditures would change by the following annual amounts and percentages:

- Private employer premiums: increase of \$52,075,000 (0.147%)
- Employee premiums by private employees, CalPERS employees, and Healthy Families enrollees: increase of \$15,150,000 (0.148%);
- Premiums for individually purchased insurance: increase of \$12,988,000 (0.340%);
- CalPERS employer premiums: increase of \$1,338,000 (0.060%);
- Medi-Cal state expenditures: increase of \$0 (0.000%)<sup>16</sup>;
- Healthy Families state expenditures: increase of \$70,000 (0.020%);
- Member copayments: increase of \$7,022,000 (0.172%) and
- Out-of-pocket payments for non-covered services: decrease of \$11,545,000 (-100%).

Individuals who currently purchase tobacco cessation services, mostly OTC medications, will realize the greatest savings under the mandate, because OTC medications would be available to them under the mandate, although they will be subject to copayments, not to exceed \$15.

These total expenditures for premiums take into account the total savings for those who quit smoking. The total savings are estimated to be \$3,418,000.<sup>17</sup> Note, these savings represent the short-term (i.e., one-year) savings resulting from reduced use of ambulatory and inpatient services among those who quit smoking, and do not account for the potential long-term savings of quitting. Long-term projections are considered beyond the scope of this analysis, but are likely to be substantial due to reductions in the rate

---

<sup>16</sup> The May 20, 2005 CHBRP analysis of SB 576 corrects certain coverage assumptions related to Medi-Cal and changes the cost estimates accordingly. This analysis reflects that Medi-Cal currently covers counseling services, prescription drugs and over-the-counter nicotine replacement therapy for smoking cessation treatment. Drugs are covered provided that the beneficiary is participating in a behavioral modification smoking cessation program.

<sup>17</sup> The cost savings calculation incorporates savings associated with reduced hospitalization for acute myocardial infarction (AMI) and savings associated with fewer low-birth weight deliveries. This analysis revises the cost savings to reflect the correct reduction in expected AMIs.



of smoking-related illnesses associated with a long lag time between exposure to smoking and onset, for example lung cancer and emphysema.

Total annual costs of smoking cessation are likely to decline in future years, as fewer smokers remain. However, cessation costs are also likely to increase in the future due to the diminishing effectiveness of smoking cessation strategies for those who continue to smoke.

#### Impact on access and health service availability (Section 3(g))

CHBRP estimates that the proposed mandate will have no impact on access to (i.e., the supply of) tobacco cessation services, because these services are already widely available and the mandate would not increase demand substantially. Expanded coverage of OTC medications would potentially encourage more insured individuals to use tobacco cessation services, but the availability of these services should not be limited by this increased demand.

### **III. PUBLIC HEALTH IMPACTS**

#### **Introduction**

The first report ever issued by a United States Surgeon General, in 1964, focused on the health effects of tobacco use. Since that time, the rate of smoking in California and nationwide has decreased dramatically, but tobacco use still remains the leading preventable cause of death and disease in the United States (DHHS, 2004). Recent estimates of smoking-related morbidity and mortality in California indicate that there were 43,137 smoking-related deaths in 1999, corresponding to 12.4 years of potential life lost, and \$131,741 in lost productivity per death (Max et al, 2004). According to the CDC, smoking results in approximately 440,000 deaths nationwide each year and generates costs of approximately \$157 billion in annual health-related economic losses. Smoking during pregnancy was estimated to result in 560 deaths in infant boys and 410 deaths in infant girls annually.

In 2004, the current Surgeon General issued the second national report on the health effects of tobacco use (DHHS, 2004). With regard to tobacco cessation, the report states that “smokers who quit can lower their risk for smoking-caused diseases and improve their health status generally. Those who never start can avoid the predictable burden of disease and lost life expectancy that results from a lifetime of smoking. Quitting smoking has immediate as well as long term benefits, reducing risks for diseases caused by smoking and improving health in general”.

During the past 15 years, Californians have had increased access to tobacco cessation education and services through the California Department of Health Services Tobacco Control Section. The California Tobacco Tax and Health Promotion Act of 1988 (Proposition 99), increased the state surtax on cigarettes and other tobacco-related products. Revenues from the “tobacco tax” were appropriated for tobacco-related research, tobacco cessation efforts and health education, and health care for medically indigent families. In 1995, California enacted a smoke-free workplace law in an effort to reduce the public health burden of second-hand smoke inhalation. During this time period, tobacco-use rates in California decreased (22% to 17.5%) and participation in cessation activities increased (48.9% to 61.5%) (Gilpin et al., 2001; Siegel et al., 2000).



By 2003, approximately 17.9% of Californians were current smokers (Table 4). Men were more likely to be current smokers than women, and those in the 18-39 year age group were slightly more likely to smoke than those aged 40-64 years. 2001 data from the California Tobacco Survey estimates that approximately two thirds of all smokers made at least one quit attempt in the prior year (Table 5). Only about a quarter of those who made a cessation attempt maintained the cessation for over 90 days. Smokers who made quit attempts used a variety of cessation assistance techniques, including counseling (7.8%), self-help materials (15.5%), and prescription and non-prescription medication (22.9%).

**Table 4.** Smoking Prevalence among California Adults (%), 2003

<i>Age (years)</i>	<i>Male</i>	<i>Female</i>	<i>Total</i>
18-24	22.7	14.1	18.5
25-39	24.0	13.4	18.7
40-64	20.2	14.2	17.1
Total	22	13.9	17.9

Source: California Health Interview Study, 2003.

**Table 5.** Tobacco Cessation Attempts

<i>Cessation Attempts</i>	<i>California Smokers, %</i>
Quit attempts*	
Quit attempt of 1 day or longer	61.5
90 or more days of abstinence	23.7
Use of cessation assistance	
Counseling	7.8
Self-help materials	15.2
Medication	22.9

Source: \*California Adult Tobacco Survey (CATS) and Behavioral Risk Factor Survey (BRFS), 1997-2001 weighted to 1990 California population. California Department of Health Services, Tobacco Control Section, 2002

Despite state-level advances in tobacco cessation, the smoking rates in California remain higher than the Healthy People 2010 target of 12% for adults, particularly for ethnic minorities and those with low socioeconomic status (Table 6). Overall, American Indians have the highest smoking rates in the state, with similar rates for men and women in this group (32%). Blacks, Whites, and those of “other” or mixed rates have similar smoking rates of approximately 20%, with women in these groups having slightly lower rates than men. Hispanics and Asians have the lowest smoking rates, with significant gender differences. Hispanic men have smoking rates that are three times higher than Hispanic women, whereas Asian men’s rates are almost four times higher than those for Asian women. Socioeconomic status also appears to play a role in smoking behavior. Both men and women with incomes less than 300% of the Federal Poverty Level are more likely to smoke than those who have higher incomes.



**Table 6.** Racial and Economic Disparities in Smoking Prevalence (%)

<i>Race</i>	<i>Male</i>	<i>Female</i>	<i>Total</i>
Hispanic	21.8	7.8	15.0
Native American	32.0	32.3	32.2
Asian	23.8	6.5	14.8
Black	22.5	18.4	20.3
White	21.2	18.0	19.6
Other/2 or more races	24.2	15.8	20.1

  

<i>Poverty Status</i>	<i>Male</i>	<i>Female</i>	<i>Total</i>
0-99% FPL	28.3	14.1	20.3
100-199% FPL	26.7	16.1	21.2
200-299% FPL	25.4	15.8	20.6
300% + FPL	18.2	12.5	15.5

FPL= Federal Poverty Level

Source: California Health Interview Study, 2003.

### Short-Term Health Outcomes

Our analysis of the potential public health impacts of SB 576 focuses on two primary outcome areas: low birth weight deliveries and acute myocardial infarction (AMI). These outcomes were chosen for analysis because (1) the medical literature shows that smoking has a direct causal relationship to these conditions, (2) there is evidence that tobacco cessation has an impact on these outcomes, and (3) there is evidence that these outcomes have the potential to develop within one year of smoking initiation, and conversely, the effects of these outcomes have the potential to be mitigated in the short term.

#### Low Birth Weight Baseline Data

A recent report from the California Department of Health Services estimated that in 1999, approximately 76.4 years of potential life were lost, per death, due to death from short gestation or low birth weight (Max et al, 2002). Low birth weight deliveries can come about either due to intrauterine growth retardation (IUGR) or pre-term birth. IUGR is a condition in which the fetus fails to grow as expected, but may be born at the appropriate time. A baby born pre-term may be growing appropriately, but be low-birth weight by virtue of the early delivery. The CDC estimates that in 1999, approximately 11.4% of pregnant women in California were smokers, and they were twice as likely as non-smokers to deliver a low birth-weight baby (12.2% vs 6.3%) (CDC, 1999). The low birthweight delivery rate as a percentage of live births in California varies by race, with Black women having significantly higher rates than women of other racial backgrounds (Table 7).



**Table 7.** Birth Outcomes: Low Birth Weight

<i>Race</i>	<i>Low birth weight delivery rate as a percentage of live births</i>
Hispanic	5.7%
White	5.9%
Black	11.7%
Native American	6.2%
Asian	7.1%
<i>Total</i>	<i>6.3%</i>

Source: March of Dimes, 2002.

### *Summary of Studies*

Many studies have shown that smoking during pregnancy increases the risk for delivery of a low birth weight baby (Messecar, 2001). According to a recent CDC report, approximately 7% of all low birth weight deliveries in California are directly attributable to smoking. The cost of low birth weight deliveries can be significant due to the increased likelihood of complications during the birth, extended hospitalization for mothers and infants, and the increased need for neonatal intensive care. A 2004 report from the CDC estimates that California's 1996 neonatal expenditures attributed to maternal smoking were \$567 per woman, relative to \$14 per non-smoker (CDC, 2004). A recent study by Adams and colleagues showed that maternal smoking increases the risk of neonatal intensive care unit admission by 20% (Adams et al., 2002).

It appears that tobacco cessation, particularly during the first trimester of pregnancy, reduces the risk of low birth weight deliveries and infant death. Salihu and colleagues estimated that 986 infant deaths could be prevented annually in the United States, if all pregnant smokers quit (Salihu et al, 2003). A 1990 study estimated that for every \$1 spent on tobacco cessation services for pregnant women, over \$3 in savings were achieved in medical care of low birth weight babies, and in the reduction of perinatal mortality (Marks et al, 1990). In 1999, Lightwood and colleagues conducted an analysis of the short-term impacts of California's public health tobacco cessation programs on the incidence of and costs associated with low birth weight deliveries. This study found that an annual 1% decrease in the smoking rate (corresponding to 3%-4% of pregnant smokers quitting) would prevent 1,300 low birth weight deliveries and save \$21 million in direct medical costs within the first year after cessation intervention.

### *Baseline population of interest*

The California Tobacco Control Section reports that approximately 11.4% of pregnant women in California are smokers. According to actuarial data from the cost and utilization section of this analysis, approximately 39,893 pregnant women smokers are currently insured in California. Of these, approximately 26,122 pregnant women smokers have coverage that includes tobacco cessation benefits and 13,771 pregnant smokers are not covered for these services. After the mandate, we expect that approximately 12,236 pregnant smokers would be newly covered for tobacco cessation services, resulting in a total of 38,348 women covered for the benefit and 1,545 remaining without coverage.



### *Expected outcome estimates without mandate*

The rate of low birth weight deliveries in California is 6.3% among non-smokers and 12.2% among smokers (CDC, 1999). In general, tobacco cessation quit rates are higher for pregnant women than for the general population of smokers. Thus, without coverage for tobacco cessation services, we would expect approximately 14% of women to quit smoking and approximately 22% of those with coverage for the benefit to quit. (Healthy People 2010 Objective 27-6), DHHS, 2004 Surgeon General's Report). Based on these assumptions, we expect that prior to the mandate, approximately 7,675 women would quit smoking during pregnancy and we would expect approximately 4,414 low birth weight deliveries.

### *Expected outcome estimates after mandate*

Under this mandate, a total of 12,236 pregnant women smokers would be newly covered for tobacco cessation benefits. We assume that a greater percentage of women would use tobacco cessation services once they become a covered benefit. If we assume that the rate of tobacco cessation for those newly covered under the mandate would increase from 14% to 22%, we would expect a total of 8,653 women to quit smoking during pregnancy under this mandate. Applying the low birth weight rate of 12.2% to the remaining smokers and 6.3% to the non-smokers, we would expect approximately 4,356 low birth weight deliveries under the mandate. Thus, we would expect 58 fewer low birth weight deliveries statewide in the year following the enactment of SB 576.

Total insured adult pregnant smokers: 39,893

Quit rate with coverage: 22%

Quit rate without coverage: 14%

Number of smokers prior to mandate: 32,218

Number of smokers after mandate: 31,240

Low birth weight rate (smokers): 12.2%

Low birth weight rate (non-smokers): 6.3%

Baseline expected cases of low birth weight deliveries: 4,414

Expected cases of low birth weight deliveries under mandate: 4,356

Total expected reduction in low birth weight deliveries due to mandate: 58

### Acute Myocardial Infarction (AMI) Baseline Data

AMI, sometimes referred to as a "heart attack," occurs when there is damage to an area of the heart due to decreased blood flow to that area. There are multiple causes of AMI, but smoking is one of the primary risk factors. Myocardial infarction is one of the leading causes of death in the United States and in California (California Center for Health Statistics). In California, the mortality rate due to AMI varies by race and gender (Table 8). For all races, men have higher AMI-related mortality than women. Black and Native American men have the highest AMI mortality rate, followed by Whites. Hispanics and Asians have the lowest mortality rates due to AMI.



**Table 8.** Acute Myocardial Infarction Mortality Rate\* by Race, California Adults Age 18-64 Years

<i>Race</i>	<i>Men</i>	<i>Women</i>
Hispanic	8.1	4.7
White	19.2	7.4
Black	26.6	20.4
Native American	23.0	10.1
Asian	13.8	3.9
<i>Total</i>	<i>16.2</i>	<i>7.2</i>

Source: California Center for Health Statistics)

Notes: \*Per 100,000 annually.

### *Summary of Studies*

The causal association between smoking and heart disease has been well documented by over 20 years of scientific evidence (Critchley and Capewell, 2003; DHHS, 2004). According to the California Department of Health Services, in 1999, men and women between the ages of 35 and 64 years who smoked were almost twice as likely to die from heart disease as were non-smokers in this age group (Max et al., 2002). According to this report, for adults between the ages of 35 and 64 years of age in 1999, there were 24.3 years of potential life lost per death from ischemic heart disease that were attributable to smoking.

A California-specific study by Lightwood and colleagues estimated the effect of the state's public health tobacco control programs on hospitalization for AMI within the first year after cessation (Lightwood and Glantz, 1997). Lightwood estimated that an annual 1% reduction in smoking prevalence (corresponding to approximately 3%-4% of smokers quitting) would result in 924 fewer hospitalizations for AMI and 538 fewer strokes which results in approximately \$44 million (1995 dollars) in savings in direct medical costs within one year. This estimate does not include deaths that occur suddenly, before transportation to a hospital can be arranged.<sup>18</sup>

Smoking is associated with both fatal and non-fatal AMI. According to recent systematic reviews of the literature on the association between smoking and heart disease, tobacco cessation is associated with a 36% reduction in risk of total mortality and a 32% reduction in risk of non-fatal AMI (Critchley and Capewell, 2003, 2004).

### *Baseline population of interest*

Approximately 2,326,775 smokers are currently insured in California. Under current coverage assumptions based on actuarial data covered in the cost and utilization section of this analysis, we expect

---

<sup>18</sup> Quantifying the short-term reduction of hospitalization as it pertains to strokes is difficult due to lack of published data on this specific outcome measure. However, Lightwood and Glantz (1997) estimated that an annual 1% reduction in smoking prevalence (corresponding to approximately 3%-4% of smokers quitting) would result in 924 fewer hospitalizations for AMI and 538 fewer hospitalizations for stroke within one year. This represents a reduction of approximately 0.6 hospitalizations for stroke for every AMI hospitalization avoided. Based on this ratio, we would expect that post-mandate, there would be approximately 9 fewer stroke-related hospitalizations statewide.



that approximately 26,983 smokers would quit smoking in any given year, resulting in 2,299,792 remaining smokers.

#### *Baseline expected outcome estimates without mandate*

Lightwood's study of the effects of California's public health tobacco cessation programs on the incidence of AMI estimates the rate of AMI in the general adult population as 0.219%. Tobacco cessation reduces the risk of AMI by approximately 32% (Critchley and Capewell, 2003, 2004), bringing the rate of AMI for non-smokers to 0.149% within the first year after cessation. According to these estimates, we would expect approximately 5077  $((26,983 \times 0.149\%) + (2,299,792 \times 0.219\%))$  baseline cases of AMI in the relevant population, prior to enactment of the mandate.

#### *Expected outcome estimates after mandate*

Based on actuarial data and assumptions listed in the cost and utilization section of this analysis, approximately 44,773 smokers would be expected to quit smoking after this mandate, resulting in 2,282,002 remaining smokers. According to these calculations, we would expect a total of 5,065  $((44,773 \times 0.149\%) + (2,282,002 \times 0.219\%))$  cases of AMI if SB 576 were enacted. Thus, under this mandate, we would expect 12 fewer cases of AMI statewide.

Total insured adult smokers: 2,326,775  
Total smokers who quit annually, prior to mandate: 26,983  
Total adult smokers prior to mandate: 2,299,792  
Total smokers who quit annually, after mandate: 44,773  
Total adult smokers after mandate: 2,282,002  
Rate of AMI among smokers 0.219%  
Rate of AMI among non-smokers 0.149%  
Baseline expected cases of AMI: 5,077  
Expected cases of AMI under mandate: 5,065  
Total expected reduction in AMI due to mandate: 12

### **Economic Costs of Smoking in California**

Smoking has both direct and indirect costs for individuals, employers, government, and society. The Surgeon General's report (DHHS, 2004) states that young smokers have poorer health status, increased work absenteeism, and use more medical care than non-smokers (e.g., 25% more health care).

Medical care makes up the largest proportion of the direct costs of smoking. The CDC reports that men who smoke incur \$15,800 (in 2002 dollars) more in lifetime medical expenses than non-smokers, and women who smoke incur \$17,500 more than non-smokers (Fellows et al, 2002). According to the California Department of Health Services, in 1999, the state spent \$8,564,623 in total health care costs directly attributable to smoking, including \$4,016,568 in hospital care, \$2,060,234 in outpatient care, and \$1,133,432 for prescriptions (Max et al, 2002). A 1995 study by Wagner and colleagues estimates that tobacco cessation resulted in significant decreases in use of outpatient and inpatient health care services (Wagner et al, 1995).



Due to the multiple effects of smoking on the body, a significant proportion of the death and disease burden of smoking will not be evident until many years after smoking is initiated. Indirect costs in terms of loss of productivity, quality of life, and life years lost are difficult to estimate for outcomes that may develop over a 30-year time period. California's Department of Health Services reports that in 1999, 12.4 years of potential life were lost due to smoking-related disease, with an associated \$5,175,678 in lost productivity for men and \$2,019,478 in lost productivity for women (Max et al, 2002). A recent study by Mulder and colleagues estimates that smokers who successfully quit report improved quality of life relative to current smokers (Mulder et al, 2001). Another study, by Taylor and colleagues, estimates the life extension achieved by tobacco cessation. Cessation at an early age (35 years old) resulted in an additional seven to eight years of life for men and an additional six to seven years of life for women. Cessation at a later age (65 years old), although resulting in significantly fewer life years gained, one to two for men and two to three for women, illustrates the benefits of cessation at any age.

### **Conclusion of public health impacts**

Based on the scientific evidence for the effectiveness of tobacco cessation services, the impact of tobacco cessation on both short-term and long-term outcomes, and the evidence of economic benefits from reductions in smoking-related expenses, SB 576 would likely have a positive impact on public health in the state. Short-term benefits would be limited to those associated with the reduction in morbidity and mortality associated with AMI and low birth weight deliveries. A discussion of long-term benefits of tobacco cessation is beyond the scope of this analysis. However, we might assume that the long-term effects of cessation would be likely to impact conditions that are more chronic in nature, possibly more disabling and more costly than the short-term outcomes discussed in this analysis.

Based on the baseline demographic and disease-related statistics for California, SB 576 could potentially have a disproportionate impact on men, ethnic minorities, and low-income individuals. This analysis has identified gender, income, and race-based disparities in smoking prevalence and rates of disease. A recent CDC report estimates that tobacco cessation rates have been increasing among non-Hispanic Blacks (Woollery et al, 2003). Presumably, a mandate that would make tobacco cessation services available without any associated fees would make these services more accessible to traditionally underserved populations. Thus we might expect to see increased utilization among these groups and presumably a greater reduction in morbidity and mortality, relative to groups with lower baseline smoking and disease rates.



## TABLES

**Table 9. Baseline (Premandate) Per Member Per Month Premium and Expenditures, California, Calendar Year 2005**

	Large Group				Small Group				Individual		Public				Total Annual Expenditures
	HMO	PPO	POS	FFS	HMO	PPO	POS	FFS	HMO	PPO	CalPERS HMO	Medi-Cal HMO Over 65	Medi-Cal HMO Other	Healthy Families HMO	Total (Members)
<b>Population currently covered</b>	7,400,000	3,220,000	457,000	19,000	1,498,000	875,000	454,000	4,000	887,000	1,065,000	795,000	354,000	2,846,000	494,000	20,368,000
<b>Average premium paid by employer</b>	\$187.97	\$283.90	\$234.95	\$240.59	\$161.28	\$234.40	\$180.93	\$181.88	\$0.00	\$0.00	\$231.96	\$235.05	\$86.12	\$58.68	\$41,860,457,000
<b>Average premium paid by employee</b>	\$50.45	\$57.87	\$51.96	\$63.25	\$83.36	\$73.27	\$94.91	\$37.09	\$214.23	\$120.38	\$44.18	\$0.00	\$0.00	\$6.32	\$14,079,831,000
<b>Total premium</b>	\$238.42	\$341.77	\$286.90	\$303.83	\$244.64	\$307.67	\$275.84	\$218.97	\$214.23	\$120.38	\$276.14	\$235.05	\$86.12	\$65.00	\$55,940,289,000
<b>Deductibles, copayments paid by members</b>	\$8.44	\$46.18	\$18.14	\$67.04	\$12.49	\$45.71	\$21.55	\$51.02	\$13.04	\$28.09	\$9.78	\$0.00	\$0.00	\$1.77	\$4,074,893,000
<b>Benefits not covered</b>	\$0.04	\$0.08	\$0.11	\$0.11	\$0.05	\$0.08	\$0.12	\$0.11	\$0.05	\$0.09	\$0.00	\$0.00	\$0.00	\$0.00	\$11,545,000
<b>Total Expenditures</b>	\$246.91	\$388.03	\$305.15	\$370.98	\$257.19	\$353.46	\$297.51	\$270.09	\$227.33	\$148.55	\$285.92	\$235.05	\$86.12	\$66.77	\$60,026,727,000

Source: California Health Benefits Review Program, 2005.

Note The population includes individuals and dependents in California who have private insurance (group and individual), or are enrolled in public plans subject to the Health and Safety Code, including CalPERS, Medi-Cal, or Healthy Families.

All population figures include enrollees aged 0-64, except the Medi-Cal population, which includes dually eligible Medicare/Medi-Cal recipients of all ages.

Employees and their dependents that receive their coverage from self-insured firms are excluded because these plans are not subject to mandates.

Key: FFS = fee for service; HMO = health maintenance organization; POS = point of service; PPO = preferred provider organization. CalPERS = California Public Employees' Retirement System.



**Table 10. Postmandate Impacts on Per Member Per Month and Total Expenditures by Insurance Plan Type, California, calendar year 2005**

	Large Group				Small Group				Individual		Public				Total Annual Expenditures
	HMO	PPO	POS	FFS	HMO	PPO	POS	FFS	HMO	PPO	CalPERS HMO	Medi-Cal HMO Over 65*	Medi-Cal HMO Other*	Healthy Families HMO	Total (Members)
<b>Population currently covered</b>	7,400,000	3,220,000	457,000	19,000	1,498,000	875,000	454,000	4,000	887,000	1,065,000	795,000	354,000	2,846,000	494,000	20,368,000
<b>Average Portion of Premium Paid by Employer</b>	\$0.2449	\$0.4064	\$0.5003	\$0.4687	\$0.2725	\$0.4109	\$0.4649	\$0.5421	\$0.0000	\$0.0000	\$0.1403	\$0.0000	\$0.0000	\$0.0117	\$53,483,000
<b>Average Portion of Premium Paid by Employee</b>	\$0.0657	\$0.0828	\$0.1106	\$0.1232	\$0.1408	\$0.1284	\$0.2439	\$0.1105	\$0.4724	\$0.6228	\$0.0267	\$0.0000	\$0.0000	\$0.0013	\$28,137,000
<b>Total Premium</b>	\$0.3107	\$0.4893	\$0.6109	\$0.5920	\$0.4133	\$0.5393	\$0.7088	\$0.6526	\$0.4724	\$0.6228	\$0.1670	\$0.0000	\$0.0000	\$0.0130	\$81,620,000
<b>Covered Benefits Paid by Member (Deductibles, copays, etc)</b>	\$0.0257	\$0.0473	\$0.0630	\$0.0603	\$0.0336	\$0.0490	\$0.0687	\$0.0624	\$0.0339	\$0.0507	\$0.0052	\$0.0000	\$0.0000	\$0.0001	\$ 7,022,000
<b>Benefits not covered</b>	-\$0.0401	-\$0.0814	-\$0.1107	-\$0.1060	-\$0.0540	-\$0.0842	-\$0.1206	-\$0.1097	-\$0.0546	-\$0.0871	\$0.0000	\$0.0000	\$0.0000	\$0.0000	\$ -11,545,000
<b>Total Expenditures</b>	\$0.2962	\$0.4553	\$0.5632	\$0.5463	\$0.3929	\$0.5041	\$0.6568	\$0.6053	\$0.4517	\$0.5864	\$0.1721	\$0.0000	\$0.0000	\$0.0131	\$ 77,097,000
<b>Percentage Impact of Mandate</b>															
<b>Insured Premiums</b>	0.130%	0.143%	0.213%	0.195%	0.169%	0.175%	0.257%	0.298%	0.221%	0.517%	0.060%	0.000%	0.000%	0.020%	0.146%
<b>Total Expenditures</b>	0.120%	0.117%	0.185%	0.147%	0.153%	0.143%	0.221%	0.224%	0.199%	0.395%	0.060%	0.000%	0.000%	0.020%	0.128%

Source: California Health Benefits Review Program, 2005.

Note The population includes individuals and dependents in California who have private insurance (group and individual), or are enrolled in public plans subject to the Health and Safety Code, including CalPERS, Medi-Cal, or Healthy Families.

All population figures include enrollees aged 0-64 years, except the Medi-Cal population, which includes dually eligible Medicare/Medi-Cal recipients of all ages.

Employees and their dependents that receive their coverage from self-insured firms are excluded because these plans are not subject to mandates.

\* The May 20 CHBRP analysis of SB 576 corrects certain coverage assumptions related to Medi-Cal and changes the cost estimates accordingly. This analysis reflects that Medi-Cal currently covers counseling services, prescription drugs and over-the-counter nicotine replacement therapy for smoking cessation treatment. Drugs are covered provided that the beneficiary is participating in a behavioral modification smoking cessation program.

Key: FFS = fee for service; HMO = health maintenance organization; POS = point of service; PPO = preferred provider organization. CalPERS = California Public Employees' Retirement System.



## APPENDIX A Literature Review Methods

SB 576 is an act to add §1367.27 to the Health and Safety Code and to add §10123.175 to the Insurance Code, relating to health coverage. SB 576 mandates coverage *for tobacco cessation services that include two courses of treatment in a 12-month period including personal counseling, which may be telephone or individual, and FDA-approved medication for tobacco cessation, including prescription and over-the-counter medications. Covered treatment shall comply with the Public Health Service sponsored 2000 clinical practice guideline, "Treating Tobacco Use and Dependence," or its successors.*

Appendix A describes the literature search for studies on the medical effectiveness of tobacco cessation interventions indicated in the mandate. The search focused on the effects of counseling, brief cessation interventions by physicians and clinical staff, and prescription and OTC tobacco cessation medications approved by the FDA on abstinence, recidivism rates, and pulmonary function. This appendix also discusses the outcomes used in the analysis of the mandate.

To “grade” the evidence for all outcome measures, the CHBRP effectiveness team uses a system<sup>19</sup> with the following categories:

1. Favorable (statistically significant effect): Findings are uniformly favorable, and many or all are statistically significant.
2. Pattern<sup>20</sup> toward favorable (but not statistically significant): Findings are generally favorable, but there may be none that are statistically significant.
3. Ambiguous/mixed evidence: Some findings are significantly favorable, and some findings with sufficient statistical power show no effect.
4. Pattern toward no effect/weak evidence: Studies generally find no effect, but this may be due to a lack of statistical power.
5. No effect: There is statistical evidence of no clinical effect in the literature with sufficient statistical power to make this assessment.
6. Unfavorable: No findings show a statistically significant benefit, and some show significant harms.
7. Insufficient evidence to make a “call”: There are very few relevant findings, so that it is difficult to discern a pattern.

The literature search was conducted in PubMed and the Cochrane Library for relevant research published between 2000 and 2005 and was limited to English abstracts. No patient populations were excluded. Publication types included in the literature search were systematic reviews, meta-analyses, randomized controlled trials, clinical trials, multicenter studies, practice guidelines, and reviews. Internet searches were conducted and focused on several Web sites, including HEDIS, CDC, U.S. Prevention Task Force, American Cancer Society, American Cancer Institute, American Lung Association, and the Task Force on Community Preventive Services.

The *Medical Subject Headings* (MeSH) terms used by the librarian in the PubMed search were:

Smoking cessation/methods

Smoking cessation/statistics & numerical data

---

<sup>19</sup> The foregoing system was adapted from the system used by the U.S. Preventive Services Task Force, available at <http://www.ahcpr.gov/clinic/3rduspstf/ratings.htm>. The medical effectiveness team also considered guidelines from the Centers for Medicare & Medicaid Services, (available at <http://www.cms.hhs.gov/mcac/8b1-i9.asp>) and guidelines from the Blue Cross and Blue Shield Association (available at <http://www.bcbs.com/tec/teccriteria.html>).

<sup>20</sup> In this instance, the word “trend” may be used synonymously with “pattern.”



Smoking/adverse effects  
Smoking/economics  
Smoking/ mortality  
Smoking/prevention & control  
Smoking/ statistics & numerical data  
Tobacco use cessation/methods  
Tobacco use disorder/therapy  
Tobacco use disorder/drug therapy  
Tobacco use/prevention & control  
Tobacco smoke pollution  
Lung neoplasms/etiology  
Lung neoplasms/mortality  
Lung neoplasms/prevention & control  
Esophageal neoplasm/etiology  
Esophageal neoplasm/mortality  
Esophageal neoplasms/prevention & control  
Pancreatic neoplasms/etiology  
Pancreatic neoplasms/prevention & control  
Pancreatic neoplasms/mortality  
Neoplasms/prevention & control  
Myocardial ischemia/etiology  
Myocardial ischemia/mortality  
Myocardial ischemia/prevention & control  
Myocardial infarction/etiology  
Myocardial infarction/mortality  
Myocardial infarction/prevention & control  
Cerebrovascular accident/etiology  
Cerebrovascular accident/mortality  
Cardiovascular diseases/ economics  
Cardiovascular diseases/etiology  
Cardiovascular diseases/mortality  
Cardiovascular diseases/ prevention & control  
Coronary disease/etiology  
Coronary disease/economics  
Coronary disease/mortality  
Coronary disease/prevention & control  
Pregnancy outcome  
Pregnancy complications/prevention & control  
Pulmonary disease, chronic obstructive/etiology  
Pulmonary disease, chronic obstructive/mortality  
Pulmonary disease, chronic  
Asthma/prevention & control  
Forced expiratory volume/physiology  
Longevity  
Quality-adjusted life years  
Mortality  
Hospitalization  
Anti-anxiety agents/ therapeutic use  
Antidepressive agents/ therapeutic use



Bupropion/therapeutic use  
Clonidine/therapeutic use  
Dopamine uptake inhibitors/ therapeutic use  
Chewing gum  
Nicotine/therapeutic use  
Nicotine/administration & dosage  
Nicotinic agonists/ administration & dosage  
Nicotinic agonists/therapeutic use  
Substance withdrawal syndrome/ drug therapy  
Drugs, non-prescription/therapeutic use  
Drugs, non-prescription/economics  
Prescriptions, drug/therapeutic use  
Prescriptions, drug/economics  
Physician's role  
Nurser's role  
Nurse practitioners  
Nursing care  
Physician assistants  
Patient education  
Counseling  
Counseling/methods  
Counseling/economics  
Directive counseling  
Behavior therapy  
Group processes  
Hotline  
Telephone  
Cost-benefit analysis  
Drug costs  
Treatment outcome  
Outcome and process assessment (health care)  
Comparative study  
Follow-up studies  
Cohort studies  
Intervention studies  
Prospective studies  
Retrospective studies  
Evidence-based medicine

Additional keywords were used to find articles (\* indicates truncation): cessation, pulmonary function test\*, FEV2/FVC, asthma, chronic obstructive pulmonary disease, COPD, angina, myocardial infarction, stroke\*, lung cancer, esophageal cancer, pancreatic cancer, cancer, second hand smok\*, emergency room visit\*, ED visit\*, brief intervention\*, individual behavioral counseling, counseling, group behavior therapy, telephone counseling, cost\*, heart attack, pregnancy outcome\*, nicotine replacement therapy, office-based system, benefit\*, physician assistant\*, nurse or nurses, nursing care, nurse practitioner\*, prescription, non-prescription, over the counter, patch, nasal spray, inhaler, lozenge, bupropion SR, bupropion, clonidine, two courses, two sessions, two attempts, 30 minutes, treatment outcome\*, smoking status, vital sign.



The literature search resulted in nine relevant meta-analyses described in Appendix B. The search resulted in 168 references, including journal articles and Web sites. This analysis does not review the results of each clinical trial but reviews the recommendations and conclusions of evidence-based reviews and meta-analyses. Many meta-analyses published in the Cochrane Library were updated in 2004 and contain articles published after the 2000 Public Health Service Clinical Guideline (PHS, 2000).

At least two reviewers screened the title and abstract of each citation returned by the literature search to determine eligibility for inclusion. Full-text articles were obtained, and reviewers reapplied the initial eligibility criteria. This analysis excluded over 150 articles because they represented a lower level of evidence or were already included in the systematic reviews and meta-analyses.

The literature search resulted in an extensive body of literature that addresses the effectiveness of the interventions described in SB 576. Recent meta-analyses and systematic reviews provide thorough analyses on the effectiveness of counseling, brief advice, and pharmacotherapy on tobacco cessation.



## APPENDIX B

### Summary of Findings on Medical Effectiveness of Tobacco Cessation Interventions

Appendix B describes the meta-analyses and evidence-based systematic reviews relevant to this analysis and summarizes their results, usually given by an odds ratio.<sup>21</sup> Table B-2 lists the results for each intervention, where the main outcome studied in these reviews is abstinence from smoking. Table 2-B also includes a “grade” for the effectiveness of each intervention (see Appendix A for a description).

**Table B-1-a. Summary of Published Studies on Effectiveness of Tobacco Cessation Interventions (Counseling and Brief Advice)**

<b>Citation</b>	<b>Type of Trial</b>	<b>Intervention vs. Comparison Group</b>	<b>Population Studied</b>	<b>Location</b>
Fiore et al., 2000	Meta-analysis	Counseling in person or via telephone in individual or group therapy, physician advice to quit vs. no advice	Smokers after five-months follow-up	N/A
Lancaster and Stead, 2004	Meta-analysis	Brief advice vs. no advice (or usual care)	Smokers after six-months follow-up	N/A
Stead, Lancaster, 2004	Meta-analysis	Group tobacco cessation program vs. no self-help materials or no intervention	Smokers after six-months follow-up	N/A
Lancaster, Stead, 2002	Meta-analysis	Face-to-face individual counseling from a health care worker not involved in routine clinical care vs. minimal intervention	Smokers after six-months follow-up	N/A
Rice and Stead, 2004	Meta-analysis	Advice by a nursing professional vs. no intervention	Adult smokers over 18 years after six-months follow-up	N/A
Rigotti et al., 2003	Meta-analysis	Intensive intervention (inpatient contact plus follow up for at least one month) vs. usual care	Hospital inpatients after six-months follow-up	N/A
Stead et al., 2003	Meta-analysis	Proactive telephone support vs. minimal intervention	Smokers after six-months follow-up	N/A

<sup>21</sup> An odds ratio can be defined as the odds of an event occurring in the intervention group divided by the odds of an event occurring in the control group. An odds ratio of 1 indicates no difference between comparison groups. For undesirable outcomes, an odd ratio of less than 1 indicates that the intervention was effective in reducing the risk of that outcome.



Citation	Type of Trial	Intervention vs. Comparison Group	Population Studied	Location
Bernstein and, Becker, 2002	Systematic review	Brief counseling (< 3 minutes counseling) vs. usual care	Emergency Department patients after six months follow-up	N/A

**Table B-1-b. Summary of Published Studies on Effectiveness of Tobacco Cessation Interventions (Medications)**

Citation	Type of Trial	Intervention vs. Comparison Group	Population Studied	Location
Fiore et al., 2000	Systematic review	Pharmacotherapy – bupriopriion SR vs. placebo, nicotine gum vs. placebo	Smokers after five-months follow-up	N/A
Silagy et al., 2004	Meta-analysis	Different forms of Nicotine Replacement therapy vs. placebo or no treatment	Smokers after six-months follow-up	N/A
Hughes et al., 2004	Meta-analysis	Medication vs. placebo or alternative therapy	Smokers after six-months follow-up	N/A

**Table B-2. Summary of Evidence of Effectiveness of Interventions on Abstinence**

**Various Types of Counseling, Favorable**

Trial	Results	Categorization of Results (Significance, Direction)
Fiore et al., 2000 (meta-analysis)	For 4-8 person-to-person treatment sessions, odds ratio for cessation = 1.9 (23 trials) Individual counseling, odds ratio = 1.7 (67 trials) Higher intensity counseling (> 10 minutes), odds ratio = 2.3 (55 trials) 91-300 minutes (total amount of contact time), odds ratio = 3.2 (16 trials)	Sig, fav Sig, fav Sig, fav Sig, fav
Stead and Lancaster, 2004 (meta-analysis, 6 trials)	Odds ratio for cessation = 2.19	Sig, fav
Lancaster and Stead, 2002 (meta-analysis)	Individual counseling from tobacco cessation specialist vs. minimal intervention, odds ratio = 1.62 (15 trials) Intensive counseling vs. brief counseling, odds ratio = 0.98 (4 trials)	Sig, fav NS, not fav



<b>Trial</b>	<b>Results</b>	<b>Categorization of Results (Significance, Direction)</b>
Rigotti et al., 2003 (meta-analysis, 6 trials)	Intensive intervention, odds ratio = 1.82	Sig, fav
Stead et al., 2003 (meta-analysis, 13 trials)	Telephone counseling, odds ratio = 1.56	Sig, fav

### **Brief Advice to Quit, Favorable**

<b>Trial</b>	<b>Results</b>	<b>Categorization of Results (Significance, Direction)</b>
Fiore et al., 2000 (meta-analysis, 7 trials)	Odds ratio = 1.3	Sig, fav
Lancaster and Stead, 2004 (meta-analysis, 17 trials)	Brief advice, odds ratio = 1.74 Intensive versus minimal advice, odds ratio = 1.44	Sig, fav
Rice and Stead, 2004 (meta-analysis, 20 trials)	Odds ratio = 1.47	Sig, fav
Bernstein and Becker, 2002 (systematic review, 16 trials)	Quit rates at 6-12 months = 3% (usual care), 8%-11% (brief counseling)	Sig, fav

### **Bupropion SR, favorable**

<b>Trial</b>	<b>Results</b>	<b>Categorization of Results (Significance, Direction)</b>
Fiore et al., 2000 (meta-analysis, 2 trials)	Efficacy on abstinence, odds ratio = 2.1	Sig, fav
Hughes et al., 2004 (meta-analysis, 19 trials)	Efficacy on abstinence, odds ratio = 2.06	Sig, fav



**Nicotine Gum, favorable**

<b>Trial</b>	<b>Results</b>	<b>Categorization of Results (Significance, Direction)</b>
Fiore et al., 2000 (meta-analysis, 13 trials)	Efficacy on abstinence, odds ratio = 1.5	Sig, fav
Silagy et al., 2004 (meta-analysis, 52 trials)	Abstinence from smoking, odds ratio = 1.66	Sig, fav

**Nicotine Nasal Spray, favorable**

<b>Trial</b>	<b>Results</b>	<b>Categorization of Results (Significance, Direction)</b>
Fiore et al., 2000 (meta-analysis, 3 trials)	Efficacy on abstinence, odds ratio = 2.7	Sig, fav
Silagy et al., 2004 (meta-analysis, 4 trials)	Abstinence from smoking, odds ratio = 2.35	Sig, fav

**Nicotine Patch, favorable**

<b>Trial</b>	<b>Results</b>	<b>Categorization of Results (Significance, Direction)</b>
Fiore et al., 2000 (meta-analysis, 27 trials)	Efficacy on abstinence, odds ratio = 1.9	Sig, fav
Silagy et al., 2004 (meta-analysis, 37 trials)	Abstinence from smoking, odds ratio = 1.81	Sig, fav



**Nicotine Inhaler, favorable**

<b>Trial</b>	<b>Results</b>	<b>Categorization of Results (Significance, Direction)</b>
Fiore et al., 2000 (meta-analysis, 4 trials)	Efficacy on abstinence, odds ratio = 2.5	Sig, fav
Silagy et al., 2004 (meta-analysis, 4 trials)	Abstinence from smoking, odds ratio = 2.14	Sig, fav

**Nicotine Lozenge, favorable**

<b>Trial</b>	<b>Results</b>	<b>Categorization of Results (Significance, Direction)</b>
Silagy et al., 2004 (meta-analysis, 4 trials)	Abstinence from smoking, odds ratio = 2.05	Sig, fav



## APPENDIX C:

### Articles Not Directly Referenced Within the Analysis or Excluded from the Analysis

Appendix C lists the articles that were either not directly referenced within this analysis or excluded from this analysis. These articles represented a lower level of evidence or were already included in the systematic reviews or meta-analyses.

Influence on lifestyle measures and five-year coronary risk by a comprehensive lifestyle intervention programme in patients with coronary heart disease. (2003).. *European Journal of Cardiovascular Prevention and Rehabilitation*. 10(6):429-437.

Bupropion to aid smoking cessation. (2000). *Drug and Therapeutics Bulletin*. 38(10):73-75.

U.S. Public Health Service. (2000). A clinical practice guideline for treating tobacco use and dependence: A US public health service report. The tobacco use and dependence clinical practice guideline panel, staff, and consortium representatives. *The Journal of the American Medical Association*. 283(24):3244-54.

Task Force on Community Preventive Services (2000). Strategies for reducing exposure to environmental tobacco smoke, increasing tobacco-use cessation, and reducing initiation in communities and health-care systems. A report on recommendations of the. *MMWR Recommendations and Report*. 49(RR-12):1-11.

Ahluwalia JS, Harris KJ, Catley D, Okuyemi KS, Mayo MS. (2002). Sustained-release bupropion for smoking cessation in african americans: A randomized controlled trial. *JAMA*.. 288(4):468-474.

Ahmadi J, Ashkani H, Ahmadi M, Ahmadi N. (2003). Twenty-four week maintenance treatment of cigarette smoking with nicotine gum, clonidine and naltrexone. *Journal of Substance Abuse and Treatment*. 24(3):251-255.

Alterman AI, Gariti P, Mulvaney F. (2001). Short- and long-term smoking cessation for three levels of intensity of behavioral treatment. *Psychology of Addictive Behaviors*. 15(3):261-264.

American Cancer Institute. *Prevention and Cessation of Cigarette Smoking: Control of Tobacco Use (pdq®)*. <http://www.cancer.gov/cancerinfo/pdq/prevention/control-of-tobacco-use/patient/>

American Lung Association. (2004). *Nicotine Replacement*.  
<http://www.lungusa.org/site/pp.asp?c=dvLUK9O0E&b=33566>

Anderson P, Jane-Llopis E. (2004). How can we increase the involvement of primary health care in the treatment of tobacco dependence? A meta-analysis. *Addiction*. 99(3):299-312.

Aubin HJ, Lebargy F, Berlin I, Bidaut-Mazel C, Chemali-Hudry J, Lagrue G. (2004). Efficacy of bupropion and predictors of successful outcome in a sample of french smokers: A randomized placebo-controlled trial. *Addiction*. 99(9):1206-1218.

Berlin I, Aubin HJ, Pedarriosse AM, Rames A, Lancrenon S, Lagrue G. (2002). Lazabemide, a selective, reversible monoamine oxidase b inhibitor, as an aid to smoking cessation. *Addiction*. 97(10):1347-1354.



- Biberman R, Neumann R, Katzir I, Gerber Y. (2003). A randomized controlled trial of oral selegiline plus nicotine skin patch compared with placebo plus nicotine skin patch for smoking cessation. *Addiction*. 98(10):1403-1407.
- Bohadana A, Nilsson F, Rasmussen T, Martinet Y. (2003). Gender differences in quit rates following smoking cessation with combination nicotine therapy: Influence of baseline smoking behavior. *Nicotine and Tobacco Research*. 5(1):111-116.
- Bolliger CT. (2000). Practical experiences in smoking reduction and cessation. *Addiction*. 95 Suppl 1:S19-24.
- Bolliger CT, Zellweger JP, Danielsson T, van Biljon X, Robidou A, Westin A, Perruchoud AP, Sawe U. (2002). Influence of long-term smoking reduction on health risk markers and quality of life. *Nicotine and Tobacco Research*. 4(4):433-439.
- Bolliger CT, Zellweger JP, Danielsson T, van Biljon X, Robidou A, Westin A, Perruchoud AP, Sawe U. (2000). Smoking reduction with oral nicotine inhalers: Double blind, randomised clinical trial of efficacy and safety. *British Medical Journal*. 321(7257):329-333.
- Borland R, Balmford J, Segan C, Livingston P, Owen N. (2003). The effectiveness of personalized smoking cessation strategies for callers to a quitline service. *Addiction*. 98(6):837-846.
- Borland R, Segan CJ, Livingston PM, Owen N. (2001). The effectiveness of callback counselling for smoking cessation: A randomized trial. *Addiction*. 96(6):881-889.
- Borrelli B, Papandonatos G, Spring B, Hitsman B and Niaura R. (2004). Experimenter-defined quit dates for smoking cessation: Adherence improves outcomes for women but not for men. *Addiction*. 99(3):378-85.
- Brandon TH, Collins BN, Juliano LM, Lavez AB. (2000). Preventing relapse among former smokers: A comparison of minimal interventions through telephone and mail. *Journal of Consulting and Clinical Psychology*. 68(1):103-113.
- Brown RA, Kahler CW, Niaura R, Abrams DB, Sales SD, Ramsey SE, Goldstein MG, Burgess ES, Miller IW. (2001). Cognitive-behavioral treatment for depression in smoking cessation. *Journal of Consulting and Clinical Psychology*. 69(3):471-480.
- Buchanan L. (2002). Implementing a smoking cessation program for pregnant women based on current clinical practice guidelines. *Journal of the American Academy of Nurse Practitioners*. 14(6):243-250.
- Buchanan LM, El-Banna M, White A, Moses S, Siedlik C, Wood M. (2004). An exploratory study of multicomponent treatment intervention for tobacco dependency. *Journal of Nursing Scholarship*. 36(4):324-330.
- Burling TA, Burling AS, Latini D. (2001). A controlled smoking cessation trial for substance-dependent inpatients. *Journal of Consulting and Clinical Psychology*. 69(2):295-304.



Carpenter MJ, Hughes JR, Solomon LJ, Callas PW. (2004). Both smoking reduction with nicotine replacement therapy and motivational advice increase future cessation among smokers unmotivated to quit. *Journal of Consulting and Clinical Psychology*. 72(3):371-81.

Centers for Disease Control and Prevention. (2005). Coverage for tobacco use cessation treatments. [http://www.cdc.gov/tobacco/educational\\_materials/cessation/ReimbursementBrochureFull.pdf](http://www.cdc.gov/tobacco/educational_materials/cessation/ReimbursementBrochureFull.pdf)

Centers for Disease Control and Prevention. (2001). Women and smoking: A report of the surgeon general 2001. [http://www.cdc.gov/tobacco/sgr/sgr\\_forwomen/index.htm](http://www.cdc.gov/tobacco/sgr/sgr_forwomen/index.htm)

Centers for Disease Control and Prevention. *Literature Review on the Effectiveness of State Tobacco Control Programs*. [http://www.cdc.gov/tobacco/sustainingstates/PreventionPrograms\\_1.htm](http://www.cdc.gov/tobacco/sustainingstates/PreventionPrograms_1.htm)

Chatkin JM, Mariante de Abreu C, Haggstrom FM, Wagner MB, Fritscher CC. (2004). Abstinence rates and predictors of outcome for smoking cessation: Do Brazilian smokers need special strategies? *Addiction*. 99(6):778-784.

Chengappa KN, Kambhampati RK, Perkins K, Nigam R, Anderson T, Brar JS, Vemulapalli HK, Atzert R, Key P, Kang JS, Levine J. (2001). Bupropion sustained release as a smoking cessation treatment in remitted depressed patients maintained on treatment with selective serotonin reuptake inhibitor antidepressants. *Journal of Clinical Psychiatry*. 62(7):503-508.

Cooper TV, DeBon MW, Stockton M, Klesges RC, Steenbergh TA, Sherrill-Mittleman D, Jennings LC, ad Johnson KC. (2004). Correlates of adherence with transdermal nicotine. *Addictive Behaviors*. 29(8):1565-1578.

Cornuz J, Humair JP, Seematter L, Stoianov R, van Melle G, Stalder H and Pecoud A. (2002). Efficacy of resident training in smoking cessation: A randomized, controlled trial of a program based on application of behavioral theory and practice with standardized patients. *Annals of Internal Medicine*. 136(6):429-437.

Covey LS, Glassman AH, Stetner F, Rivelli S, Stage K. (2002). A randomized trial of sertraline as a cessation aid for smokers with a history of major depression. *American Journal of Psychiatry*. 159(10):1731-1737.

Cox LS, Patten CA, Niaura RS, Decker PA, Rigotti N, Sachs DP, Buist AS, Hurt RD. (2004). Efficacy of bupropion for relapse prevention in smokers with and without a past history of major depression. *Journal of General Internal Medicine*. 19(8):828-834.

Croghan GA, Sloan JA, Croghan IT, Novotny P, Hurt RD, DeKrey WL, Mailliard JA, Ebbert LP, Swan DK, Walsh DJ, Wiesenfeld M, Levitt R, Stella P, Johnson PA, Tschetter LK, Loprinzi C. (2003). Comparison of nicotine patch alone versus nicotine nasal spray alone versus a combination for treating smokers: A minimal intervention, randomized multicenter trial in a nonspecialized setting. *Nicotine and Tobacco Research*. 5(2):181-187.

Curry SJ, Ludman EJ, Graham E, Stout J, Grothaus L, Lozano P. (2003). Pediatric-based smoking cessation intervention for low-income women: A randomized trial. *Archives of Pediatric and Adolescent Medicine*. 157(3):295-302.



- da Costa CL, Younes RN, Lourenco MT. (2002). Stopping smoking: A prospective, randomized, double-blind study comparing nortriptyline to placebo. *Chest*. 122(2):403-408.
- Dale LC, Ebbert JO, Schroeder DR, Croghan IT, Rasmussen DF, Trautman JA, Cox LS, Hurt RD. (2002). Bupropion for the treatment of nicotine dependence in spit tobacco users: A pilot study. *Nicotine and Tobacco Research*. 4(3):267-274.
- Dale LC, Glover ED, Sachs DP, Schroeder DR, Offord KP, Croghan IT, Hurt RD. (2001). Bupropion for smoking cessation: Predictors of successful outcome. *Chest*. 119(5):1357-1364.
- Dalsgareth OJ, Hansen NC, Soes-Petersen U, Evald T, Hoegholm A, Barber J, Vestbo J. (2004). A multicenter, randomized, double-blind, placebo-controlled, 6-month trial of bupropion hydrochloride sustained-release tablets as an aid to smoking cessation in hospital employees. *Nicotine and Tobacco Research*. 6(1):55-61.
- David S, Lancaster T, Stead LF. (2001). Opioid antagonists for smoking cessation. *Cochrane Database of Systematic Reviews*. (3):CD003086.
- Dornelas EA, Sampson RA, Gray JF, Waters D, Thompson PD. (2000). A randomized controlled trial of smoking cessation counseling after myocardial infarction. *Preventive Medicine*. 30(4):261-268.
- Dunn C, Deroo L and Rivara FP. (2001). The use of brief interventions adapted from motivational interviewing across behavioral domains: A systematic review. *Addiction*. 96(12):1725-1742.
- Durcan MJ, Johnston JA, White J, Gonzales D, Sachs DP, Rigotti N, Niaura R. (2004). Bupropion SR for relapse prevention: A "slips-allowed" analysis. *American Journal of Health Behavior*. 28(5):456-463.
- Durcan MJ, White J, Jorenby DE, Fiore MC, Rennard SI, Leischow SJ, Nides MA, Ascher JA, Johnston JA. (2002). Impact of prior nicotine replacement therapy on smoking cessation efficacy. *American Journal of Health Behavior*. 26(3):213-220.
- El-Bastawissi A, McAfee T, Zbikowski SM, Hollis J, Stark M, Wassum K, Clark N, Barwinski R, Broughton E. (2003). The uninsured and medicaid oregon tobacco user experience in a real world, phone based cessation programme. *Tobacco Control*. 12(1):45-51.
- Emmons KM, Hammond SK, Fava JL, Velicer WF, Evans JL, Monroe AD. (2001). A randomized trial to reduce passive smoke exposure in low-income households with young children. *Pediatrics*. 108(1):18-24.
- Etter JF, Laszlo E, Perneger TV. (2004). Postintervention effect of nicotine replacement therapy on smoking reduction in smokers who are unwilling to quit: Randomized trial. *Journal of Clinical Psychopharmacology*. 24(2):174-179.
- Etter JF, Laszlo E, Zellweger JP, Perrot C, Perneger TV. (2002). Nicotine replacement to reduce cigarette consumption in smokers who are unwilling to quit: A randomized trial. *Journal of Clinical Psychopharmacology*. 22(5):487-495.
- Fagerstrom KO, Hughes JR, Rasmussen T, Callas PW. (2000). Randomised trial investigating effect of a novel nicotine delivery device (eclipse) and a nicotine oral inhaler on smoking behaviour, nicotine and carbon monoxide exposure, and motivation to quit. *Tobacco Control*. 9(3):327-333.



Feeney GF, McPherson A, Connor JP, McAlister A, Young MR, Garrahy P. (2001). Randomized controlled trial of two cigarette quit programmes in coronary care patients after acute myocardial infarction. *Internal Medicine Journal*. 31(8):470-475.

Fiore MC, McCarthy DE, Jackson TC, Zehner ME, Jorenby DE, Mielke M, Smith SS, Guiliani TA, Baker TB. (2004). Integrating smoking cessation treatment into primary care: An effectiveness study. *Preventive Medicine*. 38(4):412-420.

France EK, Glasgow RE, Marcus AC. (2001). Smoking cessation interventions among hospitalized patients: What have we learned? *Preventive Medicine*. 32(4):376-388.

Garcia MP, Becona E. (2000). Evaluation of the amount of therapist contact in a smoking cessation program. *The Spanish Journal of Psychology*. 3(1):28-36.

Garcia-Vera MP. (2004). Clinical utility of the combination of cognitive-behavioral techniques with nicotine patches as a smoking-cessation treatment: Five-year results of the "ex-moker" program. *Journal of Substance Abuse Treatment*. 27(4):325-333.

Garvey AJ, Kinnunen T, Nordstrom BL, Utman CH, Doherty K, Rosner B, Vokonas PS. (2000). Effects of nicotine gum dose by level of nicotine dependence. *Nicotine and Tobacco Research*. 2(1):53-63.

George TP, Vessicchio JC, Termine A, Bregartner TA, Feingold A, Rounsaville BJ, Kosten TR. (2002). A placebo controlled trial of bupropion for smoking cessation in schizophrenia. *Biological Psychiatry*. 52(1):53-61.

George TP, Ziedonis DM, Feingold A, Pepper WT, Satterburg CA, Winkel J, Rounsaville BJ, Kosten TR. (2000). Nicotine transdermal patch and atypical antipsychotic medications for smoking cessation in schizophrenia. *American Journal of Psychiatry*. 157(11):1835-1842.

Glasgow RE, Whitlock EP, Eakin EG, Lichtenstein E. (2000). A brief smoking cessation intervention for women in low-income planned parenthood clinics. *American Journal of Public Health*. 90(5):786-789.

Glavas D, Rumboldt M, Rumboldt Z. (2003). Smoking cessation with nicotine replacement therapy among health care workers: Randomized double-blind study. *Croatian Medical Journal*. 44(2):219-224.

Glover ED, Glover PN, Franzon M, Sullivan CR, Cerullo CC, Howell RM, Keyes GG, Nilsson F, Hobbs GR. (2002). A comparison of a nicotine sublingual tablet and placebo for smoking cessation. *Nicotine and Tobacco Research*. 4(4):441-450.

Gold PB, Rubey RN, Harvey RT. (2002). Naturalistic, self-assignment comparative trial of bupropion sr, a nicotine patch, or both for smoking cessation treatment in primary care. *American Journal of Addiction*. 11(4):315-331.

Gonzales DH, Nides MA, Ferry LH, Kustra RP, Jamerson BD, Segall N, Herrero LA, Krishen A, Sweeney A, Buaron K, Metz A. (2001). Bupropion sr as an aid to smoking cessation in smokers treated previously with bupropion: A randomized placebo-controlled study. *Clinical Pharmacology and Therapeutics*. 69(6):438-444.



- Gorin SS, Heck JE. (2004). Meta-analysis of the efficacy of tobacco counseling by health care providers. *Cancer Epidemiology, Biomarkers, and Prevention*. 13(12):2012-2022.
- Gourlay SG, Stead LF, Benowitz NL. (2004). Clonidine for smoking cessation. *Cochrane Database Systematic Reviews*. (3):CD000058.
- Grandes G, Cortada JM, Arrazola A. (2000). An evidence-based programme for smoking cessation: Effectiveness in routine general practice. *British Journal of General Practice*. 50(459):803-807.
- Grandes G, Cortada JM, Arrazola A, Laka JP. (2003). Predictors of long-term outcome of a smoking cessation programme in primary care. *British Journal of General Practice*. 53(487):101-107.
- Haas AL, Munoz RF, Humfleet GL, Reus VI, Hall SM. (2004). Influences of mood, depression history, and treatment modality on outcomes in smoking cessation. *Journal of Consulting and Clinical Psychology*. 72(4):563-570.
- Hajek P, Taylor TZ, Mills P. (2002). Brief intervention during hospital admission to help patients to give up smoking after myocardial infarction and bypass surgery: Randomised controlled trial. *British Medical Journal*. 324(7329):87-89.
- Hajek P, West R, Lee A, Foulds J, Owen L, Eiser JR, Main N. (2001). Randomized controlled trial of a midwife-delivered brief smoking cessation intervention in pregnancy. *Addiction*. 96(3):485-494.
- Hall SM, Humfleet GL, Reus VI, Munoz RF, Cullen J. (2004). Extended nortriptyline and psychological treatment for cigarette smoking. *American Journal of Psychiatry*. 161(11):2100-2107.
- Hall SM, Humfleet GL, Reus VI, Munoz RF, Hartz DT, Maude-Griffin R. (2002). Psychological intervention and antidepressant treatment in smoking cessation. *Archives of General Psychiatry*. 59(10):930-936.
- Hand S, Edwards S, Campbell IA, Cannings R. (2002). Controlled trial of three weeks nicotine replacement treatment in hospital patients also given advice and support. *Thorax*. 57(8):715-718.
- Hanson K, Allen S, Jensen S, Hatsukami D. (2003). Treatment of adolescent smokers with the nicotine patch. *Nicotine and Tobacco Research*. 5(4):515-526.
- Harris KJ, Okuyemi KS, Catley D, Mayo MS, Ge B, Ahluwalia JS. (2004). Predictors of smoking cessation among african-americans enrolled in a randomized controlled trial of bupropion. *Preventive Medicine*. 38(4):498-502.
- Hatsukami DK, Rennard S, Patel MK, Kotlyar M, Malcolm R, Nides MA, Dozier G, Bars MP, Jamerson BD. (2004). Effects of sustained-release bupropion among persons interested in reducing but not quitting smoking. *American Journal of Medicine*. 116(3):151-157.
- Haug NA, Svikis DS, Diclemente C. (2004). Motivational enhancement therapy for nicotine dependence in methadone-maintained pregnant women. *Psychology of Addictive Behaviors*. 18(3):289-292.



- Hays JT, Hurt RD, Rigotti NA, Niaura R, Gonzales D, Durcan MJ, Sachs DP, Wolter TD, Buist AS, Johnston JA, White JD. (2001). Sustained-release bupropion for pharmacologic relapse prevention after smoking cessation. A randomized, controlled trial. *Annals of Internal Medicine*. 135(6):423-433.
- Hegaard HK, Kjaergaard H, Moller LF, Wachmann H, Ottesen B. (2003). Multimodal intervention raises smoking cessation rate during pregnancy. *Acta obstetrica et gynecologica Scandinavica*. 82(9):813-819.
- Hitsman B, Spring B, Borrelli B, Niaura R, Papandonatos GD. (2001). Influence of antidepressant pharmacotherapy on behavioral treatment adherence and smoking cessation outcome in a combined treatment involving fluoxetine. *Experimental and clinical psychopharmacology*. 9(4):355-362.
- Hopkins DP, Briss PA, Ricard CJ, Husten CG, Carande-Kulis VG, Fielding JE, Alao MO, McKenna JW, Sharp DJ, Harris JR, Woollery TA, Harris KW. (2001). Reviews of evidence regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. *Am J Preventive Medicine*. 20(2 Suppl):16-66.
- Hughes JR, Shiffman S, Callas P, Zhang J. (2003). A meta-analysis of the efficacy of over-the-counter nicotine replacement. *Tobacco Control*. 12(1):21-27.
- Hughes JR, Stead LF, Lancaster T. (2000). Anxiolytics and antidepressants for smoking cessation. *Cochrane Database Systematic Reviews*. (2):CD000031.
- Hughes JR, Stead LF, Lancaster T. (2000). Anxiolytics for smoking cessation. *Cochrane Database Systematic Reviews*. (4):CD002849.
- Hurt RD, Krook JE, Croghan IT, Loprinzi CL, Sloan JA, Novotny PJ, Kardinal CG, Knost JA, Tirona MT, Addo F, Morton RF, Michalak JC, Schaefer PL, Porter PA, Stella PJ. (2003). Nicotine patch therapy based on smoking rate followed by bupropion for prevention of relapse to smoking. *Journal of Clinical Oncology*. 21(5):914-920.
- Hurt RD, Wolter TD, Rigotti N, Hays JT, Niaura R, Durcan MJ, Gonzales D, Sachs DP, Johnston JA, Offord KP. (2002). Bupropion for pharmacologic relapse prevention to smoking: Predictors of outcome. *Addictive Behaviors*. 27(4):493-507.
- Jamerson BD, Nides M, Jorenby DE, Donahue R, Garrett P, Johnston JA, Fiore MC, Rennard SI, Leischow SJ. (2001). Late-term smoking cessation despite initial failure: An evaluation of bupropion sustained release, nicotine patch, combination therapy, and placebo. *Clinical Therapeutics*. 23(5):744-752.
- Johnson JL, Ratner PA, Bottorff JL, Hall W, Dahinten S. (2000). Preventing smoking relapse in postpartum women. *Nursing Research*. 49(1):44-52.
- Jones C, Griffiths RD, Skirrow P, Humphris G. (2001). Smoking cessation through comprehensive critical care. *Intensive Care Medicine*. 27(9):1547-1549.
- Joseph AM, Arikian NJ, An LC, Nugent SM, Sloan RJ, Pieper CF. (2004). Results of a randomized controlled trial of intervention to implement smoking guidelines in veterans affairs medical centers: Increased use of medications without cessation benefit. *Medical Care*. 42(11):1100-1110.



- Kennedy DT, Paulson DM, Eddy TD, Patel PC, Patkar AD, Holdford DA, Genina VY, Griffin EN. (2004). A smoking-cessation program consisting of extensive counseling, pharmacotherapy, and office spirometry: Results of a pilot project in a veterans administration medical center. *Pharmacotherapy*. 24(10):1400-1407.
- Killen JD, Fortmann SP, Schatzberg AF, Hayward C, Sussman L, Rothman M, Strausberg L, Varady A. (2000). Nicotine patch and paroxetine for smoking cessation. *Journal of Consulting and Clinical Psychology*. 68(5):883-889.
- Killen JD, Robinson TN, Ammerman S, Hayward C, Rogers J, Stone C, Samuels D, Levin SK, Green S, Schatzberg AF. (2004). Randomized clinical trial of the efficacy of bupropion combined with nicotine patch in the treatment of adolescent smokers. *Journal of Consulting and Clinical Psychology*. 72(4):729-735.
- Lancaster T, Stead LF. (2000). Self-help interventions for smoking cessation. *Cochrane Database Systematic Reviews*. (2):CD001118.
- Lancaster T, Stead LF. (2000). Mecamylamine (a nicotine antagonist) for smoking cessation. *Cochrane Database Systematic Reviews*. (2):CD001009.
- Lancaster T, Stead LF. (2000). Silver acetate for smoking cessation. *Cochrane Database Systematic Reviews*. (2):CD000191.
- Lawrence T, Aveyard P, Evans O, Cheng KK. (2003). A cluster randomised controlled trial of smoking cessation in pregnant women comparing interventions based on the transtheoretical (stages of change) model to standard care. *Tobacco Control*. 12(2):168-177.
- Leischow SJ, Ranger-Moore J, Muramoto ML, Matthews E. (2004). Effectiveness of the nicotine inhaler for smoking cessation in an otc setting. *American Journal of Health Behavior*. 28(4):291-301.
- Lerman C, Kaufmann V, Rukstalis M, Patterson F, Perkins K, Audrain-McGovern J, Benowitz N. (2004). Individualizing nicotine replacement therapy for the treatment of tobacco dependence: A randomized trial. *Annals of Internal Medicine*. 140(6):426-433.
- Lumley J, Oliver S, Waters E. (2000). Interventions for promoting smoking cessation during pregnancy. *Cochrane Database Systematic Reviews*. (2):CD001055.
- Macleod ZR, Charles MA, Arnaldi VC, Adams IM. (2003). Telephone counselling as an adjunct to nicotine patches in smoking cessation: A randomised controlled trial. *Medical Journal of Australia*. 179(7):349-352.
- Manske S, Miller S, Moyer C, Phaneuf MR, Cameron R. (2004). Best practice in group-based smoking cessation: Results of a literature review applying effectiveness, plausibility, and practicality criteria. *American Journal of Health Promotion*. 18(6):409-423.
- McAlister AL, Rabius V, Geiger A, Glynn TJ, Huang P, Todd R. (2004). Telephone assistance for smoking cessation: One year cost effectiveness estimations. *Tobacco Control*. 13(1):85-86.



- Melvin CL, Dolan-Mullen P, Windsor RA, Whiteside HP, Jr., Goldenberg RL. (2000). Recommended cessation counselling for pregnant women who smoke: A review of the evidence. *Tobacco Control*. 9 Suppl 3:III80-84.
- Molyneux A, Lewis S, Leivers U, Anderton A, Antoniak M, Brackenridge A, Nilsson F, McNeill A, West R, Moxham J, Britton J. (2003). Clinical trial comparing nicotine replacement therapy (nrt) plus brief counselling, brief counselling alone, and minimal intervention on smoking cessation in hospital inpatients. *Thorax*. 58(6):484-488.
- Munafo M, Rigotti N, Lancaster T, Stead L, Murphy M. (2001). Interventions for smoking cessation in hospitalised patients: A systematic review. *Thorax*. 56(8):656-663.
- Murray RP, Connett JE, Rand CS, Pan W, Anthonisen NR. (2002). Persistence of the effect of the lung health study (lhs) smoking intervention over eleven years. *Preventive Medicine*. 35(4):314-319.
- Murray RP, Gerald LB, Lindgren PG, Connett JE, Rand CS, Anthonisen NR. (2000). Characteristics of participants who stop smoking and sustain abstinence for 1 and 5 years in the lung health study. *Preventive Medicine*. 30(5):392-400.
- Myles PS, Leslie K, Angliss M, Mezzavia P, Lee L. (2004). Effectiveness of bupropion as an aid to stopping smoking before elective surgery: A randomised controlled trial. *Anaesthesia*. 59(11):1053-1058.
- Pbert L, Ockene JK, Zapka J, Ma Y, Goins KV, Oncken C, Stoddard AM. (2004). A community health center smoking-cessation intervention for pregnant and postpartum women. *Am J Preventive Medicine*. 26(5):377-385.
- Pelkonen M, Notkola IL, Tukiainen H, Tervahauta M, Tuomilehto J, Nissinen A. (2001). Smoking cessation, decline in pulmonary function and total mortality: A 30 year follow up study among the finnish cohorts of the seven countries study. *Thorax*. 56(9):703-707.
- Pieterse ME, Seydel ER, DeVries H, Mudde AN, Kok GJ. (2001). Effectiveness of a minimal contact smoking cessation program for Dutch general practitioners: A randomized controlled trial. *Preventive Medicine*. 32(2):182-190.
- Prochazka AV, Kick S, Steinbrunn C, Miyoshi T, Fryer GE. (2004). A randomized trial of nortriptyline combined with transdermal nicotine for smoking cessation. *Archives of Internal Medicine*. 164(20):2229-2233.
- Quist-Paulsen P, Gallefoss F. (2003). Randomised controlled trial of smoking cessation intervention after admission for coronary heart disease. *British Medical Journal*. 327(7426):1254-127.
- Rabius V, McAlister AL, Geiger A, Huang P, Todd R. (2004). Telephone counseling increases cessation rates among young adult smokers. *Health Psychology*. 23(5):539-541.
- Ratner PA, Johnson JL, Richardson CG, Bottorff JL, Moffat B, Mackay M, Fofonoff D, Kingsbury K, Miller C, Budz B. (2004). Efficacy of a smoking-cessation intervention for elective-surgical patients. *Research in Nursing and Health*. 27(3):148-161.



- Reid R, Pipe A, Higginson L, Johnson K, D'Angelo MS, Cooke D, Dafoe W. (2003). Stepped care approach to smoking cessation in patients hospitalized for coronary artery disease. *Journal of Cardiopulmonary Rehabilitation*. 23(3):176-182.
- Richards D, Toop L, Brockway K, Graham S, McSweeney B, MacLean D, Sutherland M, Parsons A. (2003). Improving the effectiveness of smoking cessation in primary care: Lessons learned. *New Zealand Medical Journal*. 116(1173):U417.
- Richman PB, Dinowitz S, Nashed AH, Eskin B, Sylvan E, Allegra C, Allegra J, Mandell M. (2000). The emergency department as a potential site for smoking cessation intervention: A randomized, controlled trial. *Academic Emergency Medicine*. 7(4):348-353.
- Schuurmans MM, Diacon AH, van Biljon X, Bolliger CT. (2004). Effect of pre-treatment with nicotine patch on withdrawal symptoms and abstinence rates in smokers subsequently quitting with the nicotine patch: A randomized controlled trial. *Addiction*. 99(5):634-640.
- Shiffman S, Di Marino ME, Pillitteri JL. (2005). The effectiveness of nicotine patch and nicotine lozenge in very heavy smokers. *Journal of Substance Abuse Treatment*. 28(1):49-55.
- Shiffman S, Dresler CM, Hajek P, Gilbert SJ, Targett DA, Strahs KR. (2002). Efficacy of a nicotine lozenge for smoking cessation. *Archives of Internal Medicine*. 162(11):1267-76.
- Shiffman S, Dresler CM, Rohay JM. (2004). Successful treatment with a nicotine lozenge of smokers with prior failure in pharmacological therapy. *Addiction*. 99(1):83-92.
- Shiffman S, Gorsline J, Gorodetzky CW. (2002). Efficacy of over-the-counter nicotine patch. *Nicotine and Tobacco Research*. 4(4):477-83.
- Shiffman S, Paty JA, Rohay JM, Di Marino ME, Gitchell JG. (2001). The efficacy of computer-tailored smoking cessation material as a supplement to nicotine patch therapy. *Drug and Alcohol Dependence*. 64(1):35-46.
- Shiffman S, Rolf CN, Hellebusch SJ, Gorsline J, Gorodetzky CW, Chiang YK, Schleusener DS, Di Marino ME. (2002). Real-world efficacy of prescription and over-the-counter nicotine replacement therapy. *Addiction*. 97(5):505-516.
- Silagy C, Stead LF. (2001). Physician advice for smoking cessation. *Cochrane Database Systematic Reviews*. (2):CD000165.
- Simon JA, Carmody TP, Hudes ES, Snyder E, Murray J. (2003). Intensive smoking cessation counseling versus minimal counseling among hospitalized smokers treated with transdermal nicotine replacement: A randomized trial. *American Journal of Medicine*. 114(7):555-562.
- Simon JA, Duncan C, Carmody TP, Hudes ES. (2004). Bupropion for smoking cessation: A randomized trial. *Archives of Internal Medicine*. 164(16):1797-1803.
- Smith PM, Reilly KR, Houston Miller N, DeBusk RF, Taylor CB. (2002). Application of a nurse-managed inpatient smoking cessation program. *Nicotine and Tobacco Research*. 4(2):211-222.



- Smith SS, Jorenby DE, Fiore MC, Anderson JE, Mielke MM, Beach KE, Piasecki TM, Baker TB. (2001). Strike while the iron is hot: Can stepped-care treatments resurrect relapsing smokers? *Journal of Consulting and Clinical Psychology*. 69(3):429-439.
- Smith SS, Jorenby DE, Leischow SJ, Nides MA, Rennard SI, Johnston JA, Jamerson B, Fiore MC, Baker TB. (2003). Targeting smokers at increased risk for relapse: Treating women and those with a history of depression. *Nicotine and Tobacco Research*. 5(1):99-109.
- Stanton WR, Lowe JB, Moffatt J, Del Mar CB. (2004). Randomised control trial of a smoking cessation intervention directed at men whose partners are pregnant. *Preventive Medicine*. 38(1):6-9.
- Stillman FA, Hartman AM, Graubard BI, Gilpin EA, Murray DM, Gibson JT. (2003). Evaluation of the american stop smoking intervention study (assist): A report of outcomes. *Journal of the National Cancer Institute*. 95(22):1681-91.
- Stotts AL, Diclemente CC, Dolan-Mullen P. (2002). One-to-one: A motivational intervention for resistant pregnant smokers. *Addictive Behaviors*. 27(2):275-92.
- Swan GE, Javitz HS, Jack LM, Curry SJ, McAfee T. (2004). Heterogeneity in 12-month outcome among female and male smokers. *Addiction*. 99(2):237-50.
- Swan GE, McAfee T, Curry SJ, Jack LM, Javitz H, Dacey S, Bergman K. (2003). Effectiveness of bupropion sustained release for smoking cessation in a health care setting: A randomized trial. *Archives of Internal Medicine*. 163(19):2337-44.
- The Task Force on Community Preventive Services. (2003). *Effectiveness of Using Provider Reminders or Provider Education, with or without Patient Education*. <http://www.thecommunityguide.org/>
- The Task Force on Community Preventive Services. (2003). *Effectiveness of Telephone Counseling and Support to Help More Tobacco Users Quit*. <http://www.thecommunityguide.org/tobacco/tobac-AJPM-evrev.pdf>
- Tonnesen P, Tonstad S, Hjalmarson A, Leborgy F, Van Spiegel PI, Hider A, Sweet R, Townsend J. (2003). A multicentre, randomized, double-blind, placebo-controlled, 1-year study of bupropion sr for smoking cessation. *Journal of Internal Medicine*. 254(2):184-192.
- Tonstad S, Farsang C, Klaene G, Lewis K, Manolis A, Perruchoud AP, Silagy C, van Spiegel PI, Astbury C, Hider A, Sweet R. (2003). Bupropion sr for smoking cessation in smokers with cardiovascular disease: A multicentre, randomised study. *European Heart Journal*. 24(10):946-955.
- U.S. Preventive Services Task Force. (2003). U.S. Preventive services task force. Counseling to prevent tobacco use and tobacco-related diseases: Recommendation statement. November 2003.
- Valanis B, Lichtenstein E, Mullooly JP, Labuhn K, Brody K, Severson HH, Stevens N. (2001). Maternal smoking cessation and relapse prevention during health care visits. *American Journal of Preventive Medicine*. 20(1):1-8.
- Wallstrom M, Nilsson F, Hirsch JM. (2000). A randomized, double-blind, placebo-controlled clinical evaluation of a nicotine sublingual tablet in smoking cessation. *Addiction*. 95(8):1161-1171.



Ward MM, Doebbeling BN, Vaughn TE, Uden-Holman T, Clarke WR, Woolson RF, Letuchy E, Branch LG, Perlin J. (2003). Effectiveness of a nationally implemented smoking cessation guideline on provider and patient practices. *Preventive Medicine*. 36(3):265-271.

Wennike P, Danielsson T, Landfeldt B, Westin A, Tonnesen P. (2003). Smoking reduction promotes smoking cessation: Results from a double blind, randomized, placebo-controlled trial of nicotine gum with 2-year follow-up. *Addiction*. 98(10):1395-1402.

West R, Hajek P, Nilsson F, Foulds J, May S, Meadows A. (2001). Individual differences in preferences for and responses to four nicotine replacement products. *Psychopharmacology*. 153(2):225-230.

Wiggers LC, Smets EM, de Haes JC, Peters RJ, Legemate DA. (2003). Smoking cessation interventions in cardiovascular patients. *European Journal of Vascular and Endovascular Surgery*. 26(5):467-475.

Wileyto P, Patterson F, Niaura R, Epstein L, Brown R, Audrain-McGovern J, Hawk L, Lerman C. (2004). Do small lapses predict relapse to smoking behavior under bupropion treatment? *Nicotine and Tobacco Research*. 6(2):357-366.

Windsor RA, Woodby LL, Miller TM, Hardin JM, Crawford MA, DiClemente CC. (2000). Effectiveness of agency for health care policy and research clinical practice guideline and patient education methods for pregnant smokers in medicaid maternity care. *American Journal of Obstetrics and Gynecology*. 182(1 Pt 1):68-75.

Wisborg K, Henriksen TB, Jespersen LB, Secher NJ. (2000). Nicotine patches for pregnant smokers: A randomized controlled study. *Obstetrics and Gynecology*. 96(6):967-971.

Zhu SH, Anderson CM, Tedeschi GJ, Rosbrook B, Johnson CE, Byrd M, Gutierrez-Terrell E. (2002). Evidence of real-world effectiveness of a telephone quitline for smokers. *New England Journal of Medicine*. 347(14):1087-1093.



## APPENDIX D

### Cost Impact Analysis: General Caveats and Assumptions

This appendix describes general caveats and assumptions used in conducting the cost impact analysis. For additional information on the cost model and underlying methodology, please refer to the CHBRP Web site, <http://www.chbrp.org/costimpact.html>.

The cost analysis in this report was prepared by Milliman and University of California, Los Angeles (UCLA), with the assistance of CHBRP staff. Per the provisions of AB 1996 (California Health and Safety Code, §127660, et seq.), the analysis includes input and data from an independent actuarial firm, Milliman. In preparing cost estimates, Milliman and UCLA relied on a variety of external data sources. The *Milliman Health Cost Guidelines* (HCG) were used to augment the specific data gathered for this mandate. The HCGs are updated annually and are widely used in the health insurance industry to estimate the impact of plan changes on health care costs. Although the data were reviewed for reasonableness, they were used without independent audit.

The expected costs in this report are not predictions of future costs. Instead, they are estimates of the costs that would result if a certain set of assumptions were exactly realized. Actual costs will differ from these estimates for a wide variety of reasons, including:

- Prevalence of mandated benefits before and after the mandate different from our assumptions.
- Utilization of mandated services before and after the mandate different from our assumptions.
- Random fluctuations in the utilization and cost of health care services.

Additional assumptions that underlie the cost estimates presented here are:

- Cost impacts are only shown for people with insurance.
- The projections do not include people covered under self-insurance employer plans because those employee benefit plans are not subject to state-mandated minimum benefit requirements.
- Employers and employees will share proportionately (on a percentage basis) in premium rate increases resulting from the mandate. In other words, the distribution of premium paid by the subscriber (or employee) and the employer will be unaffected by the mandate.

There are other variables that may affect costs, but which Milliman did not consider in the cost projections presented in this report. Such variables include, but are not limited to:

- Population shifts by type of health insurance coverage. If a mandate increases health insurance costs, then some employer groups or individuals may elect to drop their coverage. Employers may also switch to self-funding to avoid having to comply with the mandate.
- Changes in benefit plans. To help offset the premium increase resulting from a mandate, members or insured may elect to increase their overall plan deductibles or copayments. Such changes would have a direct impact on the distribution of costs between the health plan and the insured person, and may also result in utilization reductions (i.e., high levels of patient cost sharing result in lower utilization of health care services). Milliman did not include the effects of such potential benefit changes in its analysis.
- Adverse Selection. Theoretically, individuals or employer groups who had previously foregone insurance may now elect to enroll in an insurance plan postmandate because they perceive that it is to their economic benefit to do so.



- Health plans may react to the mandate by tightening their medical management of the mandated benefit. This would tend to dampen our cost estimates. The dampening would be more pronounced on the plan types that previously had the least restrictive medical management (i.e., FFS and PPO plans).
- Variation in existing utilization and costs, and in the impact of the mandate, by geographic area and delivery system models: Even within the plan types we modeled (HMO, PPO, POS, and FFS), there are variations in utilization and costs within California. One source of difference is geographic. Utilization differs within California due to differences in the health status of the local commercial population, provider practice patterns, and the level of managed care available in each community. The average cost per service would also vary due to different underlying cost levels experienced by providers throughout California and the market dynamic in negotiations between health plans and providers.

Both the baseline costs prior to the mandate and the estimated cost impact of the mandate could vary within the state due to geographic and delivery system differences. For purposes of this analysis, however, we have estimated the impact on a statewide level.



**APPENDIX E**  
**Information Submitted by Outside Parties for Consideration for CHBRP Analysis**

In accordance with CHBRP policy to analyze information submitted by outside parties during the first two weeks of the CHBRP review, the following parties chose to submit information.

*American Cancer Society*

Support letter for SB 576 from James K. Knox, Vice President, Legislative Advocacy  
March 2, 2005

*California Tobacco Control Alliance*

Information and articles submitted on March 7, 2005, including:

Zhu SH, Anderson CM, Tedeschi GJ, Rosbrook B, Johnson CE, Byrd M, Gutierrez-Terrell E. (2002). Evidence of real-world effectiveness of a telephone quitline for smokers. *The New England Journal of Medicine*. 347(14):1087-1093.

Zhu SH, Tedeschi GJ, Anderson CM, Rosbrook B, Byrd M, Johnson CE, Gutierrez-Terrell E. (2000). Telephone counseling as adjuvant treatment for nicotine replacement therapy in a real-world setting. *Preventative Medicine*. 31: 357-363

Zhu SH, Stretch V, Balabanis M, Rosbrook B, Sadler G, Pierce J, (1996). Telephone counseling for smoking cessation: Effects of single-session and multiple session interventions. *Journal of Consulting and Clinical Psychology*. 64(1): 202-211.

*Center for Tobacco Cessation*

Support letter for SB 576 from Sara Hutchinson, Manager, Federal and State Relations  
March 2, 2005

*Cathy McDonald, MD, MPH*

Alameda County Alcohol, Tobacco and Other Drug Treatment Provider Network  
Support letter for SB 576, March 1, 2005

*William V. Corr, Executive Director, National Center for Tobacco-Free Kids*

Support letter for SB 576, March 2, 2005

*Paul Knepprath, Vice President, American Lung Association of California*

Support letter for SB 576, March 1, 2005

*Chris Kotsen, Psy.D.*, Tobacco Dependence Treatment Specialist, Coordinator,  
Tobacco Quitcenter, Behavior Health Outpatient Services, Somerset Medical Center  
Hillsborough, New Jersey

Support letter for SB 576 dated March 7, 2005

For information on the processes for submitting information to CHBRP for review and consideration please visit: <http://www.chbrp.org/requests.html>



## REFERENCES

- Adams EK, Miller VP, Ernst C, Nishimura BK, Melvin C, Merritt R. (2002). Neonatal health care costs related to smoking during pregnancy. *Health Economics*. 11(3):193-206.
- Bernstein SL, Becker BM. (2002). Preventive care in the emergency department: Diagnosis and management of smoking and smoking-related illness in the emergency department: A systematic review. *Academic Emergency Medicine*. 9(7):720-729.
- Centers for Disease Control and Prevention (CDC). (2001). *Investment in Tobacco Control: State Highlights—2001*. Atlanta, GA: Centers for Disease Control, Office on Smoking and Health.
- Centers for Disease Control and Prevention (CDC). (2004). State estimates of neonatal health care costs associated with maternal smoking—United States, 1996. *MMWR*. 53(39):915-917.
- Centers for Disease Control and Prevention (CDC). (2005). *Prenatal Smoking Databook 1999*. Atlanta, GA: Centers for Disease Control and Prevention.  
<http://www.cdc.gov/reproductivehealth/PrenatalSmkbk/index.htm> (accessed 9 April 2005).
- Centers for Medicare and Medicaid Services (CMS). (2005). *Decision Memo for Smoking and Tobacco Use Cessation Counseling (CAG-00241N)*. <http://www.cms.hhs.gov/mcd/viewdecisionmemo.asp?id=130> (accessed March 28, 2005).
- Critchley JA, Capewell S. (2003). Mortality risk reduction associated with smoking cessation in patients with coronary heart disease: A systematic review. *JAMA*. 290(1):86-97.
- Critchley J, Capewell S. (2004). Smoking cessation for the secondary prevention of coronary heart disease. *Cochrane Database of Systematic Reviews*. (1):CD003041.
- Curry SJ, Grothaus LC, McAfee T, Pabiniak C. (1998). Use and cost effectiveness of smoking-cessation services under four insurance plans in a health maintenance organization. *The New England Journal of Medicine*. 339(10):673-679.
- Fellows JL, Trosclair A., Adams EK, Rivera CC. (2002). Annual smoking-attributable mortality, years of potential life lost, and economic costs—United States, 1995–1999. *MMWR*. 51(14):300-303.
- Fichtenberg CM, Glantz SA. (2000). Association of the California Tobacco Control Program with declines in cigarette consumption and mortality from heart disease. *The New England Journal of Medicine*. 343(24): 1772-1777.
- Fiore MC, Bailey WC, Cohen SJ, et al. (2000). *Treating Tobacco Use and Dependence*. Clinical Practice Guideline. Rockville, MD: U.S. Department of Health and Human Services. Public Health Service.
- Javitz HS, Swan GE, Zbikowski SM, Curry SJ, McAfee TA, Decker D, Patterson R, Jack LM. (2004). Return on investment of different combinations of bupropion sr dose and behavioral treatment for smoking cessation in a health care setting: An employer's perspective. *Value in Health*. 7(5):535-543.



Gilpin EA, Emery SL, Farkas AJ, Distefan JM, White MM, Pierce JP, (2001). *The California Tobacco Control Program: A Decade of Progress, Results from the California Tobacco Survey, 1990-1999*. La Jolla, CA: University of California, San Diego.

Hughes J, Stead L, Lancaster T. (2004). Antidepressants for smoking cessation. *Cochrane Database Systematic Reviews*. (4):CD000031.

Kaper J, Wagena EJ, Severens JL, Van Schayck CP. (2005). Healthcare financing systems for increasing the use of tobacco dependence treatment. *The Cochrane Database of Systematic Reviews*. (1): CD004305

Katz DA, Muehlenbruch DR, Brown RB, Fiore MC, Baker TB. (2002). Effectiveness of a clinic-based strategy for implementing the AHRQ Smoking Cessation Guideline in primary care. *Preventive Medicine*. 35(3):293-301.

Katz DA, Muehlenbruch DR, Brown RL, Fiore MC, Baker TB; AHRQ Smoking Cessation Guideline Study Group. (2004). Effectiveness of implementing the Agency for Healthcare Research and Quality smoking cessation clinical practice guideline: A randomized, controlled trial. *Journal of the National Cancer Institute*. 96(8):594-603.

Lancaster T, Stead LF. (2002). Individual behavioural counselling for smoking cessation. *Cochrane Database Systematic Reviews*. (3):CD001292.

Lancaster T, Stead L. (2004). Physician advice for smoking cessation. *Cochrane Database Systematic Reviews*. (4):CD000165.

Lawrence D, Graber JE, Mills SL, Meissner HI, Warnecke R. (2003). Smoking cessation interventions in U.S. racial/ethnic minority populations: An assessment of the literature. *Preventive Medicine*. 36(2):204-216.

Legislative Analyst's Office. (2002). *2002-0: Overview of the Governor's Budget*. [http://www.lao.ca.gov/2002/budget\\_overview/2002-03\\_bud\\_overview.pdf](http://www.lao.ca.gov/2002/budget_overview/2002-03_bud_overview.pdf) (accessed 9 April 2005)

Li CQ, Windsor RA, Hassan M. (1994). Cost differences between low birthweight attributable to smoking and low birthweight for all causes. *Preventive Medicine*. 23(1):28-34.

Lightwood J. (2003). The economics of smoking and cardiovascular disease. *Progress in Cardiovascular Diseases*. 46(1): 39-78.

Lightwood J, Fleischmann KE, Glanz SA. (2001). Smoking cessation in heart failure: It is never too late. *The Journal of the American College of Cardiology*. 37(6):1683-1684.

Lightwood JM, Glantz SA. (1997). Short-term economic and health benefits of smoking cessation: Myocardial infarction and stroke. *Circulation*. 96(4):1089-1096.

Lightwood JM, Phibbs CS, Glanz SA. (1999). Short-term health and economic benefits of smoking cessation: low birth weight. *Pediatrics*. 104(6):1312-1320.

Marks JS, Koplan JP, Hogue CJ, Dalmat ME. (1990). A cost-benefit/cost-effectiveness analysis of smoking cessation for pregnant women. *American Journal Preventive Medicine*. 6(5): 282-289.



- Martin JA, Hamilton BE, Sutton PD, Ventura SJ, Menacker F, Munson ML. (2003). Births: Final data for 2002. *National Vital Statistics Reports*. 52(10):1-113.
- Max W, Rice DP, Sung HY, Zhang X, Miller L. (2004). The economic burden of smoking in California. *Tobacco Control*. 13(3):264-267.
- McGovern PG, Pankow JS, Shahar E, Doliszny KM, Folsom AR, Blackburn H, Luepker RV. (1996). Recent trends in acute coronary heart disease—mortality, morbidity, medical care, and risk factors. The Minnesota Heart Survey Investigators. *The New England Journal of Medicine*. 334(14):884-890.
- Messecar DC. (2001). Smoking cessation interventions for pregnant women to prevent low birth weight: What does the evidence show? *Journal of the American Academy of Nurse Practitioners*. 13(4):171-177.
- Mojica WA, Suttorp MJ, Sherman SE, Morton SC, Roth EA, Maglione MA, Rhodes SL, Shekelle PG. (2004). Smoking-cessation interventions by type of provider: A meta-analysis. *American Journal of Preventive Medicine*. 26(5):391-401.
- Mulder I, Tjhuis M, Smit HA, Kromhout D. (2001). Smoking cessation and quality of life: The effect of amount of smoking and time since quitting. *Preventive Medicine*. 33(6):653-660.
- Phares TM, Morrow B, Lansky A, Barfield WD, Prince CB, Marchi KS, Braveman PA, Williams LM, Kinniburgh B. (2004). Surveillance for disparities in maternal health-related behaviors—selected states, Pregnancy Risk Assessment Monitoring System (PRAMS), 2000-2001. *MMWR Surveillance Summaries*. 53(4): 1-13.
- Rice VH, Stead LF. (2004). Nursing interventions for smoking cessation. *Cochrane Database Systematic Reviews*. (1):CD001188.
- Rigotti NA, Munafo MR, Murphy MF, Stead LF. (2003). Interventions for smoking cessation in hospitalised patients. *Cochrane Database Systematic Reviews*. (1):CD001837.
- Ringen K, Anderson N, McAfee T, Zbikowski SM, Fales D. (2002). Smoking cessation in a blue-collar population: Results from an evidence-based pilot program. *American Journal of Industrial Medicine*. 42(5):367-377.
- Salihu HM, Aliyu MH, Pierre-Louis BJ, Alexander GR. (2003). Levels of excess infant deaths attributable to maternal smoking during pregnancy in the United States. *Maternal and Child Health Journal*. 7(4):219-227.
- Schauffler HH, McMenamin S, Olson K, Boyce-Smith G, Rideout JA, Kamil J. (2001). Variations in treatment benefits influence smoking cessation: Results of a randomised controlled trial. *Tobacco Control*. 10(2):175-180.
- Siegel M, Mowery PD, Pechacek TP, Strauss WJ, Schooley MW, Merritt RK, Novotny TE, Giovino GA, Eriksen MP. (2000). Trends in adult cigarette smoking in California compared with the rest of the United States, 1978-1994. *American Journal of Public Health*. 90(3):372-379.



Silagy C, Lancaster T, Stead L, Mant D, Fowler G. (2004). Nicotine replacement therapy for smoking cessation. *Cochrane Database Systematic Reviews*. (3):CD000146.

Stead LF, Lancaster T. (2004). Group behaviour therapy programmes for smoking cessation. *Cochrane Database of Systematic Reviews*. (3):CD001007.

Stead LF, Lancaster T, Perera R. (2003). Telephone counselling for smoking cessation. *Cochrane Database Systematic Reviews*. (1):CD002850.

U.S. Department of Health and Human Services (DHHS). (2000). *Evidence Report and Evidence-Based Recommendations: Interventions to Promote Smoking Cessation in the Medicare Population*. Rockville, MD: Center for Medicaid and Medicare Services.

U.S. Department of Health and Human Services (DHHS). (2004). *The Health Consequences of Smoking: A Report of the Surgeon General*. Washington, DC: Centers for Disease Control and Prevention, Office on Smoking and Health.

U.S. Public Health Service (PHS). (2000). *Treating Tobacco Use and Dependence*. Clinical Practice Guideline. Rockville, MD: U.S. Department of Health and Human Services. Public Health Service.

Wagner EH, Curry SJ, Grothaus L, Saunders KW, McBride CM. (1995). The impact of smoking and quitting on health care use. *Archives of Internal Medicine*. 155(16):1789-1795.

Woollery T, Trosclair A., Husten C, Caraballo RC, Kahende J. (2003). Cigarette smoking among adults—United States, 2001. *Morbidity and Mortality Weekly Report*. 52(40):953-956.



## California Health Benefits Review Program Committees and Staff

A group of faculty and staff undertakes most of the analysis that informs reports by the California Health Benefits Review Program (CHBRP). The CHBRP **Faculty Task Force** comprises rotating representatives from six University of California (UC) campuses and three private universities in California. In addition to these representatives, there are other ongoing contributors to CHBRP from UC. This larger group provides advice to the CHBRP staff on the overall administration of the program and conducts much of the analysis. The CHBRP **staff** coordinates the efforts of the Faculty Task Force, works with Task Force members in preparing parts of the analysis, and coordinates all external communications, including those with the California Legislature. The level of involvement of members of CHBRP's Faculty Task Force and staff varies on each report, with individual participants more closely involved in the preparation of some reports and less involved in others.

As required by CHBRP's authorizing legislation, UC contracts with a certified actuary, Milliman, to assist in assessing the financial impact of each benefit mandate bill. Milliman also helped with the initial development of CHBRP's methods for assessing that impact.

The **National Advisory Council** provides expert reviews of draft analyses and offers general guidance on the program to CHBRP staff and the Faculty Task Force. CHBRP is grateful for the valuable assistance and thoughtful critiques provided by the members of the National Advisory Council. However, the Council does not necessarily approve or disapprove of or endorse this report. CHBRP assumes full responsibility for the report and the accuracy of its contents.

### Faculty Task Force

**Helen Halpin, PhD\***, *Vice Chair for Public Health Impacts*, University of California, Berkeley  
**Gerald Kominski, PhD**, *Vice Chair for Financial Impacts*, University of California, Los Angeles  
**Edward Yelin, PhD**, *Vice Chair for Medical Effectiveness*, University of California, San Francisco  
**Wayne S. Dysinger, MD, MPH**, Loma Linda University Medical Center  
**Theodore Ganiats, MD**, University of California, San Diego  
**Sheldon Greenfield, MD**, University of California, Irvine  
**Richard Kravitz, MD**, University of California, Davis  
**Thomas MaCurdy, PhD**, Stanford University  
**Thomas Valente, PhD**, University of Southern California

### Other Contributors

**Wade Aubry, MD**, University of California, San Francisco  
**Yali Bair, PhD**, University of California, Davis  
**Nicole Bellows, MHSA**, University of California, Berkeley  
**Patricia Franks, BA**, University of California, San Francisco  
**Janet Keyzer, RN-C, MPA**, University of California, Davis  
**Christina Kuenneth, MPH**, University of California, Davis  
**Harold Luft, PhD**, University of California, San Francisco  
**Miriam Laugesen, PhD\***, University of California, Los Angeles  
**Noelle Lee, BA**, University of California, San Francisco  
**Sara McMenamin, PhD\***, University of California, Berkeley  
**Nadereh Pourat, PhD**, University of California, Los Angeles

\* Professors Halpin, Laugesen, and McMenamin recused themselves from participation or review of this analysis



## National Advisory Council

**Susan Dentzer**, Health Correspondent, *News Hour with Jim Lehrer*, PBS, Alexandria, Virginia, *Chair*

**John Bertko, FSA, MAAA**, Vice President and Chief Actuary, Humana, Inc., Louisville, KY

**Deborah Chollet, PhD**, Senior Fellow, Mathematica Policy Research, Washington, DC

**Michael Connelly, JD**, President and CEO, Catholic Healthcare Partners, Cincinnati, OH

**Maureen Cotter, ASA**, Founder, Maureen Cotter & Associates, Inc., Dearborn, MI

**Patricia Danzon, PhD**, Celia Z. Moh Professor, The Wharton School, University of Pennsylvania, Philadelphia, PA

**Joseph Ditre, JD**, Executive Director, Consumers for Affordable Health Care, Augusta, ME

**Jack Ebeler, MPA**, President and CEO, Alliance of Community Health Plans, Washington, DC

**Allen D. Feezor**, Chief Planning Officer, University Health System of Eastern Carolina, Greenville, NC

**Charles Chip Kahn, MPH**, President and CEO, Federation of American Hospitals, Washington, DC

**Lauren LeRoy, PhD**, President and CEO, Grantmakers In Health, Washington, DC

**Trudy Lieberman**, Health Policy Editor, Consumers Union, Yonkers, NY

**Devidas Menon, PhD, MHSA**, Executive Director and CEO, Institute of Health Economics, Edmonton, AB

**Marilyn Moon, PhD**, Vice President and Director, Health Program, American Institutes for Research,  
Silver Spring, MD

**Michael Pollard, JD, MPH**, Consultant, Federal Policy and Regulation, Medco Health Solutions, Washington, DC

**Karen Pollitz**, Project Director, Georgetown University Health Policy Institute, Washington, DC

**Christopher Queram**, Chief Executive Officer, Employer Health Care Alliance Cooperative, Madison, WI

**Richard Roberts, MD, JD**, Professor of Family Medicine, University of Wisconsin-Madison, Madison, WI

**Frank Samuel, LLB**, Science and Technology Advisor, Governor's Office, State of Ohio, Columbus, OH

**Roberto Tapia-Conyer, MD, MPH, MSc**, Senior Professor, National University of Mexico, Cuauhtémoc, Mexico

**Prentiss Taylor, MD**, Vice President, Medical Affairs, Amerigroup, Chicago, IL

**Reed V. Tuckson, MD**, Senior Vice President, UnitedHealth Care, Minnetonka, MN

**Judith Wagner, PhD**, Scholar-in-Residence, Institute of Medicine, Washington, DC

**Dale Whitney**, Corporate Health and Welfare Manager, UPS, Atlanta, GA

**Ronald A. Williams**, President, Aetna, Inc., Hartford, CT

## CHBRP Staff

**Michael E. Gluck, PhD**, Director

**Susan Phillip, MPP**

Assistant Director

**Sharon Culpepper**

Assistant to the Director,

CHBRP Administrative Assistant

**Sachin Kumar, BA**

Analyst

**Robert O'Reilly, BS**

Consultant

**Cynthia Robinson, MPP**

Principal Analyst

**California Health Benefits Review Program**

**1111 Franklin Street, 11<sup>th</sup> Floor**

**Oakland, CA 94607**

**Tel: 510-287-3876 Fax: 510-987-9715**

**info@chbrp.org www.chbrp.org**

The California Health Benefits Review Program is administered by the Division of Health Affairs at the University of California Office of the President, Cathryn L. Nation, Interim Vice President.

